

Recombinant

DGRmAb®

**COX IV (DGR11126) Rabbit mAb**

db11070

Package : 10µL 20µL 100µL 500µL 1mL

**Product Name** : COX IV (DGR11126) Rabbit mAb**Cat.No.:** db11070**Synonyms** : COX4; COXIV; COX4-1; COXIV-1; COX IV-1**Application** : WB, IHC-P, ICC/IF, FC, IP**Reactivity** : Human,Mouse,Rat**Host species** : Rabbit**Background**

Cytochrome c oxidase (COX) is the terminal enzyme of the mitochondrial respiratory chain. It is a multi-subunit enzyme complex that couples the transfer of electrons from cytochrome c to molecular oxygen and contributes to a proton electrochemical gradient across the inner mitochondrial membrane. The complex consists of 13 mitochondrial- and nuclear-encoded subunits. The mitochondrially-encoded subunits perform the electron transfer and proton pumping activities. The functions of the nuclear-encoded subunits are unknown but they may play a role in the regulation and assembly of the complex. This gene encodes the nuclear-encoded subunit IV isoform 1 of the human mitochondrial respiratory chain enzyme. It is located at the 3' of the NOC4 (neighbor of COX4) gene in a head-to-head orientation, and shares a promoter with it. Pseudogenes related to this gene are located on chromosomes 13 and 14. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Jan 2016]

**Immunogen**

A synthetic peptide of human COX IV

**Gene ID**

1327

**Swiss Prot**

P13073

**Synonyms**

COX4; COXIV; COX4-1; COXIV-1; COX IV-1

**Reactivity**

Human,Mouse,Rat

**Application**

WB, IHC-P, ICC/IF, FC, IP

**Recommended dilution**

WB: 1:1000

IHC-P: 1:200-1:500

ICC/IF: 1:500-1:1000


FC: 1:20

IP: 1:20


**Calculated MW**

20 kDa

Observed MW	17 kDa
Host species	Rabbit
Clonality	Monoclonal
Clonality No.	DGR11126
Isotype	IgG
Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.



Western blot analysis of extracts from Mouse heart, Rat heart tissue using db11070 at 1:1000.



Immunofluorescence analysis of HepG2 cells labelling COX IV with db11070.

The cells were fixed with 4% PFA (10min, RT) followed by treatment with 0.1% Triton X-100 (10min, RT), and blocked in 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween 20 for 1h. The cells were then incubate with db11070 (1:500) at room temprature for 1h, followed by a further incubation at room temperature for 45min with Goat Anti Rabbit IgG (H+L)-AF488 (db10005, shown in green). Nuclear DNA was labeled in blue with DAPI.

Control: Secondary antibody only.