







Smad2/3 (DGR11226) Rabbit mAb

db11146 Package : 10μL 20μL 50μL 100μL

Product Name: Smad2/3 (DGR11226) Rabbit mAb

Cat.No.: db11146

Synonyms: JV18; MADH2; MADR2; JV18-1; hMAD-2; hSMAD2

Application: WB, ICC/IF, FC, IP **Reactivity**: Human, Mouse, Rat

Host species: Rabbit

Background

The protein encoded by this gene belongs to the SMAD, a family of proteins similar to the gene products of the Drosophila gene 'mothers against decapentaplegic' (Mad) and the C. elegans gene Sma. SMAD proteins are signal transducers and transcriptional modulators that mediate multiple signaling pathways. This protein mediates the signal of the transforming growth factor (TGF)-beta, and thus regulates multiple cellular processes, such as cell proliferation, apoptosis, and differentiation. This protein is recruited to the TGF-beta receptors through its interaction with the SMAD anchor for receptor activation (SARA) protein. In response to TGF-beta signal, this protein is phosphorylated by the TGF-beta receptors. The phosphorylation induces the dissociation of this protein with SARA and the association with the family member SMAD4. The association with SMAD4 is important for the translocation of this protein into the nucleus, where it binds to target promoters and forms a transcription repressor complex with other cofactors. This protein can also be phosphorylated by activin type 1 receptor kinase, and mediates the signal from the activin. Alternatively spliced transcript variants have been observed for this gene. [provided by RefSeq,

May 2012]

Immunogen Recombinant protein of human Smad2

Gene ID 4087, 4088

Swiss Prot Q15796, P84022

Synonyms JV18; MADH2; MADR2; JV18-1; hMAD-2; hSMAD2

Reactivity Human, Mouse, Rat

Application WB, ICC/IF, FC, IP

Recommended dilution WB: 1:1000

ICC/IF: 1:100-1:200 FC: 1:200-1:500 IP: 1:20-1:50





Calculated MW 52,48 kDa

Observed MW 58,62 kDa

Host species Rabbit

Clonality Monoclonal

Clonality No. DGR11226

Isotype IgG


Purity Affinity Purification

Conjugation Un-conjugated

Storage Stability Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium


azide and 0.05% BSA. Stable for 12 months from date of receipt.

Western blot analysis of extracts from 293 cells using db11146 at 1:1000.



20 -

15 -



Western blot analysis of extracts from HeLa, MCF-7 cells using db11146 at 1:1000.