







## Cyclin D1 (DGR11113) Rabbit mAb

db11659 Package : 10μL 20μL 50μL 100μL

Product Name: Cyclin D1 (DGR11113) Rabbit mAb

Cat.No.: db11659

**Synonyms:** BCL1; PRAD1; U21B31; D11S287E

**Application :** WB, IHC-P, ICC/IF, IP **Reactivity :** Human, Mouse, Rat

Host species: Rabbit

**Background** The protein encoded by this gene belongs to the highly conserved cyclin family, whose members

are characterized by a dramatic periodicity in protein abundance throughout the cell cycle. Cyclins function as regulators of CDK kinases. Different cyclins exhibit distinct expression and degradation patterns which contribute to the temporal coordination of each mitotic event. This cyclin forms a complex with and functions as a regulatory subunit of CDK4 or CDK6, whose activity is required for cell cycle G1/S transition. This protein has been shown to interact with tumor suppressor protein Rb

and the expression of this gene is regulated positively by Rb. Mutations, amplification and

overexpression of this gene, which alters cell cycle progression, are observed frequently in a variety

of tumors and may contribute to tumorigenesis. [provided by RefSeq, Jul 2008]

**Immunogen** A synthetic peptide of human Cyclin D1

Gene ID 595

Swiss Prot P24385

Synonyms BCL1; PRAD1; U21B31; D11S287E

Reactivity Human, Mouse, Rat

**Application** WB, IHC-P, ICC/IF, IP

Recommended dilution WB: 1:2000-1:20000

IHC-P: 1:100-1:500

ICC/IF: 1:50 IP: 1:20-1:50

34 kDa

Observed MW 36 kDa

Calculated MW

Host species Rabbit



## For Research Use Only **Product Datasheet**

**Clonality** Monoclonal

Clonality No. DGR11113

**Isotype** IgG

**Purity** Affinity Purification

**Conjugation** Un-conjugated

Storage Stability Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium

azide and 0.05% BSA. Stable for 12 months from date of receipt.