

Recombinant

DGRmAb®

TriMethyl-Histone H3 (Lys4) (DGR15420) Rabbit mAb

db11677

Package : 10µL 20µL 50µL 100µL

Product Name : TriMethyl-Histone H3 (Lys4) (DGR15420) Rabbit mAb**Cat.No.:** db11677**Synonyms :** H3/j; H3C1; H3C2; H3C3; H3C4; H3C6; H3C7; H3C8; H3FJ; H3C10; H3C11; HIST1H3J**Application :** WB, ICC/IF, FC, IP, ChIP**Reactivity :** Human, Mouse, Rat**Host species :** Rabbit**Background**

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails but instead contain a palindromic termination element. This gene is found in the small histone gene cluster on chromosome 6p22-p21.3. [provided by RefSeq, Aug 2015]

Immunogen

A synthetic methyl-peptide corresponding to residues surrounding Lys4 of human Histone H3

Gene ID

8356

Swiss Prot

P68431

Synonyms

H3/j; H3C1; H3C2; H3C3; H3C4; H3C6; H3C7; H3C8; H3FJ; H3C10; H3C11; HIST1H3J

Reactivity

Human, Mouse, Rat

Application

WB, ICC/IF, FC, IP, ChIP

Recommended dilutionWB: 1:1000
ICC/IF: 1:200-1:500
FC: 1:200-1:500
IP: 1:20**Calculated MW**

15 kDa


Observed MW

17 kDa

Host species

Rabbit

Clonality	Monoclonal
Clonality No.	DGR15420
Isotype	IgG
Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.



Western blot analysis of extracts from HeLa cells using db11677 at 1:1000.