Recombinant DGRmAb®


PTEN (DGR14385) Rabbit mAb

db12505 **Package : 10μL 20μL 50μL 100μL**

Product Name : PTEN (DGR14385) Rabbit mAb
Cat.No.: db12505
Synonyms : BZS; DEC; CWS1; GLM2; MHAM; TEP1; MMAC1; PTEN1; 10q23del; PTENbeta
Application : WB, IHC-P, FC
Reactivity : Human,Mouse,Rat
Host species : Rabbit

Background	This gene was identified as a tumor suppressor that is mutated in a large number of cancers at high frequency. The protein encoded by this gene is a phosphatidylinositol-3,4,5-trisphosphate 3-phosphatase. It contains a tensin like domain as well as a catalytic domain similar to that of the dual specificity protein tyrosine phosphatases. Unlike most of the protein tyrosine phosphatases, this protein preferentially dephosphorylates phosphoinositide substrates. It negatively regulates intracellular levels of phosphatidylinositol-3,4,5-trisphosphate in cells and functions as a tumor suppressor by negatively regulating AKT/PKB signaling pathway. The use of a non-canonical (CUG) upstream initiation site produces a longer isoform that initiates translation with a leucine, and is thought to be preferentially associated with the mitochondrial inner membrane. This longer isoform may help regulate energy metabolism in the mitochondria. A pseudogene of this gene is found on chromosome 9. Alternative splicing and the use of multiple translation start codons results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Feb 2015]
Immunogen	Recombinant protein of human PTEN
Gene ID	5728
Swiss Prot	P60484
Synonyms	BZS; DEC; CWS1; GLM2; MHAM; TEP1; MMAC1; PTEN1; 10q23del; PTENbeta
Reactivity	Human,Mouse,Rat
Application	WB, IHC-P, FC
Recommended dilution	WB: 1:1000-1:5000 IHC-P: 1:50-1:100 FC: 1:100
Calculated MW	47 kDa
Observed MW	54 kDa

Host species	Rabbit
Clonality	Monoclonal
Clonality No.	DGR14385
Isotype	IgG
Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.



Western blot analysis of extracts from HeLa cells using db12505 at 1:1000.