

Recombinant

DGRmAb®

## Eph receptor B4/HTK (DGR33743) Rabbit mAb

db13057

Package : 10µL 20µL 50µL 100µL

**Product Name** : Eph receptor B4/HTK (DGR33743) Rabbit mAb**Cat.No.:** db13057**Synonyms** : HTK; MYK1; HFASD; CMAVM2; LMPHM7; TYRO11**Application** : WB, IP**Reactivity** : Human**Host species** : Rabbit**Background**

Ephrin receptors and their ligands, the ephrins, mediate numerous developmental processes, particularly in the nervous system. Based on their structures and sequence relationships, ephrins are divided into the ephrin-A (EFNA) class, which are anchored to the membrane by a glycosylphosphatidylinositol linkage, and the ephrin-B (EFNB) class, which are transmembrane proteins. The Eph family of receptors are divided into 2 groups based on the similarity of their extracellular domain sequences and their affinities for binding ephrin-A and ephrin-B ligands. Ephrin receptors make up the largest subgroup of the receptor tyrosine kinase (RTK) family. The protein encoded by this gene binds to ephrin-B2 and plays an essential role in vascular development. [provided by RefSeq, Jul 2008]

**Immunogen**

A synthetic peptide of human Eph receptor B4/HTK

**Gene ID**

2050

**Swiss Prot**

P54760

**Synonyms**

HTK; MYK1; HFASD; CMAVM2; LMPHM7; TYRO11

**Reactivity**

Human

**Application**

WB, IP

**Recommended dilution**WB: 1:1000  
IP: 1:20-1:50**Calculated MW**

108 kDa

**Observed MW**

135 kDa

**Host species**

Rabbit

**Clonality**

Monoclonal

**Clonality No.**

DGR33743

|                          |   |
|--------------------------|---|
| <b>Isotype</b>           | IgG   |
| <b>Purity</b>            | Affinity Purification   |
| <b>Conjugation</b>       | Un-conjugated   |
| <b>Storage Stability</b> | Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt. |

Western blot analysis of extracts from HepG2 cells using db13057 at 1:1000.

