Recombinant DGRmAb®


Fibronectin (DGR33303) Rabbit mAb

db13198 **Package : 10μL 20μL 50μL 100μL**


Product Name : Fibronectin (DGR33303) Rabbit mAb
Cat.No.: db13198
Synonyms : FN; CIG; FNZ; MSF; ED-B; FINC; GFND; LETS; GFND2; SMDCF
Application : WB, IHC-P, ICC/IF
Reactivity : Human,Mouse,Rat
Host species : Rabbit

Background	This gene encodes fibronectin, a glycoprotein present in a soluble dimeric form in plasma, and in a dimeric or multimeric form at the cell surface and in extracellular matrix. The encoded preproprotein is proteolytically processed to generate the mature protein. Fibronectin is involved in cell adhesion and migration processes including embryogenesis, wound healing, blood coagulation, host defense, and metastasis. The gene has three regions subject to alternative splicing, with the potential to produce 20 different transcript variants, at least one of which encodes an isoform that undergoes proteolytic processing. The full-length nature of some variants has not been determined. [provided by RefSeq, Jan 2016]
Immunogen	A synthetic peptide of human Fibronectin
Gene ID	2335
Swiss Prot	P02751
Synonyms	FN; CIG; FNZ; MSF; ED-B; FINC; GFND; LETS; GFND2; SMDCF
Reactivity	Human,Mouse,Rat
Application	WB, IHC-P, ICC/IF
Recommended dilution	WB: 1:1000 IHC-P: 1:200-1:2000 ICC/IF: 1:50
Calculated MW	263 kDa
Observed MW	263 kDa
Host species	Rabbit
Clonality	Monoclonal
Clonality No.	DGR33303

Isotype	IgG
Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.



Western blot analysis of extracts from Human plasma tissue using db13198 at 1:1000.



Immunofluorescence analysis of HeLa cells labelling Fibronectin with db13198.

The cells were fixed with cold 100% methanol (10min, 4℃) and blocked in 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween 20 for 1h. The cells were then incubate with db13198 (1:50) at room temprature for 1h, followed by a further incubation at room temperature for 45min with Goat Anti Rabbit IgG (H+L)-AF488 (db10005, shown in green). Nuclear DNA was labeled in blue with DAPI.

Control: Secondary antibody only.