Recombinant DGRmAb®

# Phospho-FADD (Ser194) (DGR16654) Rabbit mAb

**db13926** **Package : 10μL 20μL 50μL 100μL**

**Product Name :** Phospho-FADD (Ser194) (DGR16654) Rabbit mAb
**Cat.No.:** db13926
**Synonyms :** GIG3; MORT1
**Application :** WB, IP
**Reactivity :** Human
**Host species :** Rabbit

| | |
|---|---|
| **Background** | The protein encoded by this gene is an adaptor molecule that interacts with various cell surface receptors and mediates cell apoptotic signals. Through its C-terminal death domain, this protein can be recruited by TNFRSF6/Fas-receptor, tumor necrosis factor receptor, TNFRSF25, and TNFSF10/TRAIL-receptor, and thus it participates in the death signaling initiated by these receptors. Interaction of this protein with the receptors unmasks the N-terminal effector domain of this protein, which allows it to recruit caspase-8, and thereby activate the cysteine protease cascade. Knockout studies in mice also suggest the importance of this protein in early T cell development. [provided by RefSeq, Jul 2008] |
| **Immunogen** | A synthetic phosphopeptide corresponding to residues surrounding Ser194 of human FADD |
| **Gene ID** | 8772 |
| **Swiss Prot** | Q13158 |
| **Synonyms** | GIG3; MORT1 |
| **Reactivity** | Human |
| **Application** | WB, IP |
| **Recommended dilution** | WB: 1:1000<br>IP: 1:20 |
| **Calculated MW** | 23 kDa |
| **Observed MW** | 28 kDa |
| **Host species** | Rabbit |
| **Clonality** | Monoclonal |
| **Clonality No.** | DGR16654 |

| | |
|---|---|
| **Isotype** | IgG |
| **Purity** | Affinity Purification |
| **Conjugation** | Un-conjugated |
| **Storage Stability** | Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt. |