

Recombinant

DGRmAb®

53BP1 (DGR20780) Rabbit mAb

db14226

Package : 10µL 20µL 50µL 100µL

Product Name : 53BP1 (DGR20780) Rabbit mAb**Cat.No.:** db14226**Synonyms** : TP53; p202; 53BP1; TDRD30; p53BP1**Application** : WB, IHC-P, ICC/IF**Reactivity** : Human**Host species** : Rabbit**Background**

This gene encodes a protein that functions in the DNA double-strand break repair pathway choice, promoting non-homologous end joining (NHEJ) pathways, and limiting homologous recombination. This protein plays multiple roles in the DNA damage response, including promoting checkpoint signaling following DNA damage, acting as a scaffold for recruitment of DNA damage response proteins to damaged chromatin, and promoting NHEJ pathways by limiting end resection following a double-strand break. These roles are also important during V(D)J recombination, class switch recombination and at unprotected telomeres. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Aug 2017]

Immunogen

A synthetic peptide of human 53BP1

Gene ID

7158

Swiss Prot

Q12888

Synonyms

TP53; p202; 53BP1; TDRD30; p53BP1

Reactivity

Human

Application

WB, IHC-P, ICC/IF

Recommended dilution

WB: 1:5000-1:50000

IHC-P: 1:50-1:100

ICC/IF: 1:100-1:200

Calculated MW

214 kDa

Observed MW

450 kDa

Host species

Rabbit


Clonality

Monoclonal


Clonality No.

DGR20780

Isotype	IgG
Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.



Western blot analysis of extracts from HeLa cells using db14226 at 1:1000.



Immunofluorescence analysis of HepG2 cells labelling 53BP1 with db14226.

The cells were fixed with 4% PFA (10min, RT) followed by treatment with 0.1% Triton X-100 (10min, RT), and blocked in 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween 20 for 1h. The cells were then incubate with db14226 (1:100) at room temprature for 1h, followed by a further incubation at room temperature for 45min with Goat Anti Rabbit IgG (H+L)-AF488 (db10005, shown in green). Nuclear DNA was labeled in blue with DAPI.

Control: Secondary antibody only.