

Recombinant

DGRmAb®

Cytokeratin 17 (DGR19639) Rabbit mAb

db14474

Package : 10µL 20µL 50µL 100µL

Product Name : Cytokeratin 17 (DGR19639) Rabbit mAb

Cat.No.: db14474

Synonyms : PC; K17; PC2; 39.1; CK-17; PCHC1

Application : WB, IHC-P, ICC/IF, FC

Reactivity : Human,Mouse,Rat

Host species : Rabbit


Background	This gene encodes the type I intermediate filament chain keratin 17, expressed in nail bed, hair follicle, sebaceous glands, and other epidermal appendages. Mutations in this gene lead to Jackson-Lawler type pachyonychia congenita and steatocystoma multiplex. [provided by RefSeq, Aug 2008]
Immunogen	A synthetic peptide of human Cytokeratin 17
Gene ID	3872
Swiss Prot	Q04695
Synonyms	PC; K17; PC2; 39.1; CK-17; PCHC1
Reactivity	Human,Mouse,Rat
Application	WB, IHC-P, ICC/IF, FC
Recommended dilution	WB: 1:1000-1:5000 IHC-P: 1:100-1:200 ICC/IF: 1:200-1:500 FC: 1:10-1:100
Calculated MW	48 kDa
Observed MW	48 kDa
Host species	Rabbit
Clonality	Monoclonal
Clonality No.	DGR19639
Isotype	IgG
Purity	Affinity Purification

Conjugation


Un-conjugated

Storage Stability

Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.



Western blot analysis of extracts from SCC-9 cells using db14474 at 1:1000.



Immunofluorescence analysis of HeLa cells labelling Cytokeratin 17 with db14474.

The cells were fixed with 4% PFA (10min, RT) followed by treatment with 0.1% Triton X-100 (10min, RT), and blocked in 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween 20 for 1h. The cells were then incubate with db14474 (1:200) at room temperature for 1h, followed by a further incubation at room temperature for 45min with Goat Anti Rabbit IgG (H+L)-AF488 (db10005, shown in green). Nuclear DNA was labeled in blue with DAPI.

Control: Secondary antibody only.