







TRAF2 (DGR19365) Rabbit mAb

db14520 Package : 10μL 20μL 50μL 100μL

Product Name: TRAF2 (DGR19365) Rabbit mAb

Cat.No.: db14520

Synonyms: TRAP; TRAP3; MGC:45012

Application: WB, IHC-P, ICC/IF

Reactivity : Human

Host species : Rabbit

Background The protein encoded by this gene is a member of the TNF receptor associated factor (TRAF)

protein family. TRAF proteins associate with, and mediate the signal transduction from members of the TNF receptor superfamily. This protein directly interacts with TNF receptors, and forms a heterodimeric complex with TRAF1. This protein is required for TNF-alpha-mediated activation of MAPK8/JNK and NF-kappaB. The protein complex formed by this protein and TRAF1 interacts with the inhibitor-of-apoptosis proteins (IAPs), and functions as a mediator of the anti-apoptotic signals from TNF receptors. The interaction of this protein with TRADD, a TNF receptor associated apoptotic signal transducer, ensures the recruitment of IAPs for the direct inhibition of caspase activation. BIRC2/c-IAP1, an apoptosis inhibitor possessing ubiquitin ligase activity, can unbiquitinate and induce the degradation of this protein, and thus potentiate TNF-induced apoptosis. Multiple alternatively spliced transcript variants have been found for this gene, but the biological validity of only one transcript has been determined. [provided by RefSeq, Jul 2008]

Immunogen Recombinant protein of human TRAF2

Gene ID 7186

Swiss Prot Q12933

Synonyms TRAP; TRAP3; MGC:45012

Reactivity Human

Application WB, IHC-P, ICC/IF

Recommended dilution WB: 1:1000-1:5000

IHC-P: 1:50-1:100 ICC/IF: 1:100-1:200

56 kDa

Observed MW 53 kDa

Calculated MW



For Research Use Only **Product Datasheet**

Host species Rabbit

Clonality Monoclonal

Clonality No. DGR19365

Isotype IgG

Purity Affinity Purification

Conjugation Un-conjugated

Storage Stability Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium

azide and 0.05% BSA. Stable for 12 months from date of receipt.

Western blot analysis of extracts from MCF-7 cells using db14520 at 1:1000.