

Recombinant

DGRmAb®

CD13 (DGR18974) Rabbit mAb

db14591

Package : 10µL 20µL 50µL 100µL

Product Name : CD13 (DGR18974) Rabbit mAb**Cat.No.:** db14591**Synonyms** : APN; CD13; LAP1; P150; PEPN; GP150**Application** : WB, IHC-P, ICC/IF**Reactivity** : Human,Mouse,Rat**Host species** : Rabbit**Background**

Aminopeptidase N is located in the small-intestinal and renal microvillar membrane, and also in other plasma membranes. In the small intestine aminopeptidase N plays a role in the final digestion of peptides generated from hydrolysis of proteins by gastric and pancreatic proteases. Its function in proximal tubular epithelial cells and other cell types is less clear. The large extracellular carboxyterminal domain contains a pentapeptide consensus sequence characteristic of members of the zinc-binding metalloproteinase superfamily. Sequence comparisons with known enzymes of this class showed that CD13 and aminopeptidase N are identical. The latter enzyme was thought to be involved in the metabolism of regulatory peptides by diverse cell types, including small intestinal and renal tubular epithelial cells, macrophages, granulocytes, and synaptic membranes from the CNS. Human aminopeptidase N is a receptor for one strain of human coronavirus that is an important cause of upper respiratory tract infections. Defects in this gene appear to be a cause of various types of leukemia or lymphoma. [provided by RefSeq, Jul 2008]

Immunogen

A synthetic peptide of human CD13

Gene ID

290

Swiss Prot

P15144

Synonyms

APN; CD13; LAP1; P150; PEPN; GP150

Reactivity

Human,Mouse,Rat

Application

WB, IHC-P, ICC/IF

Recommended dilution

WB: 1:1000-1:5000

IHC-P: 1:200-1:2000

ICC/IF: 1:100


Calculated MW

110 kDa


Observed MW

160 kDa

Host species	Rabbit
Clonality	Monoclonal
Clonality No.	DGR18974
Isotype	IgG
Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.



Western blot analysis of extracts from THP-1 cells using db14591 at 1:1000.



Western blot analysis of extracts from HepG2 cells using db14591 at 1:1000.