

Recombinant

DGRmAb®

## TRAF2 (DGR15352) Rabbit mAb

db15082

Package : 10µL 20µL 50µL 100µL

**Product Name :** TRAF2 (DGR15352) Rabbit mAb**Cat.No.:** db15082**Synonyms :** TRAP; TRAP3; MGC:45012**Application :** WB, IHC-P, ICC/IF, FC, IP**Reactivity :** Human,Mouse,Rat**Host species :** Rabbit**Background**

The protein encoded by this gene is a member of the TNF receptor associated factor (TRAF) protein family. TRAF proteins associate with, and mediate the signal transduction from members of the TNF receptor superfamily. This protein directly interacts with TNF receptors, and forms a heterodimeric complex with TRAF1. This protein is required for TNF-alpha-mediated activation of MAPK8/JNK and NF-kappaB. The protein complex formed by this protein and TRAF1 interacts with the inhibitor-of-apoptosis proteins (IAPs), and functions as a mediator of the anti-apoptotic signals from TNF receptors. The interaction of this protein with TRADD, a TNF receptor associated apoptotic signal transducer, ensures the recruitment of IAPs for the direct inhibition of caspase activation. BIRC2/c-IAP1, an apoptosis inhibitor possessing ubiquitin ligase activity, can ubiquitinate and induce the degradation of this protein, and thus potentiate TNF-induced apoptosis. Multiple alternatively spliced transcript variants have been found for this gene, but the biological validity of only one transcript has been determined. [provided by RefSeq, Jul 2008]

**Immunogen**

A synthetic peptide of human TRAF2

**Gene ID**

7186

**Swiss Prot**

Q12933

**Synonyms**

TRAP; TRAP3; MGC:45012

**Reactivity**

Human,Mouse,Rat

**Application**

WB, IHC-P, ICC/IF, FC, IP

**Recommended dilution**

WB: 1:1000

IHC-P: 1:100-1:200

ICC/IF: 1:100


FC: 1:100

IP: 1:20-1:50


**Calculated MW**

56 kDa

Observed MW	53 kDa
Host species	Rabbit
Clonality	Monoclonal
Clonality No.	DGR15352
Isotype	IgG
Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.



Western blot analysis of extracts from THP-1, HEK293T, HeLa cells using db15082 at 1:1000.



Western blot analysis of extracts from MOLT-4 cells using db15082 at 1:1000.