

Recombinant

5 Lipoxygenase (DGR14745) Rabbit mAb

DGRmAb[®]

db15240

Package : 10µL 20µL 50µL 100µL

Product Name : 5 Lipoxygenase (DGR14745) Rabbit mAb Cat.No.: db15240 Synonyms : 5-LO; 5LPG; LOG5; 5-LOX Application : WB, IHC-P, ICC/IF, FC Reactivity : Human,Rat Host species : Rabbit

Background	This gene encodes a member of the lipoxygenase gene family and plays a dual role in the
	synthesis of leukotrienes from arachidonic acid. The encoded protein, which is expressed
	specifically in bone marrow-derived cells, catalyzes the conversion of arachidonic acid to 5(S)-
	hydroperoxy-6-trans-8,11,14-cis-eicosatetraenoic acid, and further to the allylic epoxide 5(S)-trans-
	7,9-trans-11,14-cis-eicosatetrenoic acid (leukotriene A4). Leukotrienes are important mediators of
	a number of inflammatory and allergic conditions. Mutations in the promoter region of this gene
	lead to a diminished response to antileukotriene drugs used in the treatment of asthma and may
	also be associated with atherosclerosis and several cancers. Alternatively spliced transcript
	variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jan 2012]
Immunogen	A synthetic peptide of human 5 Lipoxygenase
Gene ID	240
Swiss Prot	P09917
Synonyms	5-LO; 5LPG; LOG5; 5-LOX
Reactivity	Human,Rat
Application	WB, IHC-P, ICC/IF, FC
Recommended dilution	WB: 1:1000
	IHC-P: 1:100
	ICC/IF: 1:100
	FC: 1:100
Calculated MW	78 kDa
Observed MW	78 kDa
Host species	Rabbit

dvagbvo 戴格生物

Clonality	Monoclonal
Clonality No.	DGR14745
lsotype	lgG
Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.
K562	Western blot analysis of extracts from K562 cells using db15240 at 1:1000.
kDa 180 - 130 -	
100 - 70 - ── ←	
55 -	
40 -	
35 -	
25 -	