

Recombinant**DGRmAb®****AIF (DGR13883) Rabbit mAb****db15482****Package : 10µL 20µL 50µL 100µL****Product Name :** AIF (DGR13883) Rabbit mAb**Cat.No.:** db15482**Synonyms :** AIF; CMT2D; CMTX4; COWCK; DFNX5; NADMR; NAMSD; PDCD8; COXP6**Application :** WB, IHC, ICC/IF, FC, IP**Reactivity :** Human, Mouse, Rat**Host species :** Rabbit**Background**

This gene encodes a flavoprotein essential for nuclear disassembly in apoptotic cells, and it is found in the mitochondrial intermembrane space in healthy cells. Induction of apoptosis results in the translocation of this protein to the nucleus where it affects chromosome condensation and fragmentation. In addition, this gene product induces mitochondria to release the apoptogenic proteins cytochrome c and caspase-9. Mutations in this gene cause combined oxidative phosphorylation deficiency 6 (COXP6), a severe mitochondrial encephalomyopathy, as well as Cowchock syndrome, also known as X-linked recessive Charcot-Marie-Tooth disease-4 (CMTX-4), a disorder resulting in neuropathy, and axonal and motor-sensory defects with deafness and cognitive disability. Alternative splicing results in multiple transcript variants. A related pseudogene has been identified on chromosome 10. [provided by RefSeq, Aug 2015]

Immunogen

A synthetic peptide of human AIF

Gene ID

9131

Swiss Prot

O95831

Synonyms

AIF; CMT2D; CMTX4; COWCK; DFNX5; NADMR; NAMSD; PDCD8; COXP6

Reactivity

Human, Mouse, Rat

Application

WB, IHC, ICC/IF, FC, IP

Recommended dilution

WB: 1:1000

IHC: 1:100

ICC/IF: 1:100

FC: 1:20

IP: 1:20


Calculated MW

67 kDa

Observed MW

67 kDa

Host species	Rabbit
Clonality	Monoclonal
Clonality No.	DGR13883
Isotype	IgG
Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.



Immunofluorescence analysis of HeLa cells labelling AIF with db15482.

The cells were fixed with 4% PFA (10min, RT) followed by treatment with 0.1% Triton X-100 (10min, RT), and blocked in 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween 20 for 1h. The cells were then incubate with db15482 (1:100) at room temprature for 1h, followed by a further incubation at room temperature for 45min with Goat Anti Rabbit IgG (H+L)-AF488 ([db10005](#), shown in green). Nuclear DNA was labeled in blue with DAPI.

Control: Secondary antibody only.