

Recombinant

DGRmAb®

Urokinase (DGR13804) Rabbit mAb

db15494

Package : 10µL 20µL 50µL 100µL

Product Name : Urokinase (DGR13804) Rabbit mAb**Cat.No.:** db15494**Synonyms** : ATF; QPD; UPA; URK; u-PA; BDPLT5**Application** : WB, IHC-P**Reactivity** : Human,Mouse,Rat**Host species** : Rabbit**Background**

This gene encodes a secreted serine protease that converts plasminogen to plasmin. The encoded preproprotein is proteolytically processed to generate A and B polypeptide chains. These chains associate via a single disulfide bond to form the catalytically inactive high molecular weight urokinase-type plasminogen activator (HMW-uPA). HMW-uPA can be further processed into the catalytically active low molecular weight urokinase-type plasminogen activator (LMW-uPA). This low molecular weight form does not bind to the urokinase-type plasminogen activator receptor. Mutations in this gene may be associated with Quebec platelet disorder and late-onset Alzheimer's disease. Alternative splicing results in multiple transcript variants, at least one of which encodes an isoform that is proteolytically processed. [provided by RefSeq, Jan 2016]

Immunogen

A synthetic peptide of human Urokinase

Gene ID

5328

Swiss Prot

P00749

Synonyms

ATF; QPD; UPA; URK; u-PA; BDPLT5

Reactivity

Human,Mouse,Rat

Application

WB, IHC-P

Recommended dilutionWB: 1:1000
IHC-P: 1:100-1:200**Calculated MW**

49 kDa

Observed MW

49 kDa

Host species

Rabbit

Clonality

Monoclonal

Clonality No.

DGR13804

Isotype	IgG
Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.