

**DGRmAb**<sup>®</sup>

Recombinant

## SMURF 2 (DGR13586) Rabbit mAb

db15537

Package : 10µL 20µL 50µL 100µL

Product Name : SMURF 2 (DGR13586) Rabbit mAb Cat.No.: db15537 Synonyms : SMURF2 Application : WB Reactivity : Human,Mouse,Rat Host species : Rabbit

Background	

E3 ubiquitin-protein ligase which accepts ubiquitin from an E2 ubiquitin-conjugating enzyme in the form of a thioester and then directly transfers the ubiquitin to targeted substrates. Interacts with SMAD1 and SMAD7 in order to trigger their ubiquitination and proteasome-dependent degradation. In addition, interaction with SMAD7 activates autocatalytic degradation, which is prevented by interaction with SCYE1. Forms a stable complex with the TGF-beta receptor-mediated phosphorylated SMAD2 and SMAD3. In this way, SMAD2 may recruit substrates, such as SNON, for ubiquitin-mediated degradation. Enhances the inhibitory activity of SMAD7 and reduces the transcriptional activity of SMAD2. Coexpression of SMURF2 with SMAD1 results in considerable decrease in steady-state level of SMAD1 protein and a smaller decrease of SMAD2 level.

Immunogen	A synthetic peptide of human SMURF 2
mmunogen	

Gene ID	64750
Swiss Prot	Q9HAU4

Synonyms SMURF2

Reactivity Human,Mouse,Rat

Application WB

Recommended dilutionWB: 1:1000Calculated MW86 kDa

Observed MW 86 kDa

Host species Rabbit

Clonality Monoclonal DGR13586

www.diagbio.com

## dvagbvo 戴格生物

lsotype	lgG
Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.
kDa <sup>C2C12</sup> 140 -	Western blot analysis of extracts from C2C12 cells using db15537 at 1:1000.
98 - ← 63 -	
49 - 39 - 34 - 22 - 20 -	

15 -10 -