

Recombinant

DGRmAb®

## cIAP2 (DGR13172) Rabbit mAb

db15636

Package : 10µL 20µL 50µL 100µL

**Product Name** : cIAP2 (DGR13172) Rabbit mAb**Cat.No.:** db15636**Synonyms** : AIP1; API2; MIHC; CIAP2; HAIP1; HIAP1; MALT2; RNF49; c-IAP2**Application** : WB, IHC-P, FC, IP**Reactivity** : Human**Host species** : Rabbit**Background**

This gene encodes a member of the IAP family of proteins that inhibit apoptosis by binding to tumor necrosis factor receptor-associated factors TRAF1 and TRAF2, probably by interfering with activation of ICE-like proteases. The encoded protein inhibits apoptosis induced by serum deprivation but does not affect apoptosis resulting from exposure to menadione, a potent inducer of free radicals. It contains 3 baculovirus IAP repeats and a ring finger domain. Transcript variants encoding the same isoform have been identified. [provided by RefSeq, Aug 2011]

**Immunogen**

A synthetic peptide of human cIAP2

**Gene ID**

330

**Swiss Prot**

Q13489

**Synonyms**

AIP1; API2; MIHC; CIAP2; HAIP1; HIAP1; MALT2; RNF49; c-IAP2

**Reactivity**

Human

**Application**

WB, IHC-P, FC, IP

**Recommended dilution**

WB: 1:2000-1:20000

IHC-P: 1:200-1:500

FC: 1:50-1:100

IP: 1:20-1:50

**Calculated MW**

68 kDa

**Observed MW**

72 kDa

**Host species**

Rabbit

**Clonality**

Monoclonal

**Clonality No.**

DGR13172

Isotype	IgG
Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.



Western blot analysis of extracts from Raji cells using db15636 at 1:1000.