

Recombinant

DGRmAb®

Cyclin D2 (DGR12228) Rabbit mAb

db15797

Package : 10µL 20µL 50µL 100µL

Product Name : Cyclin D2 (DGR12228) Rabbit mAb**Cat.No.:** db15797**Synonyms** : MPPH3; KIAK0002**Application** : WB, ICC/IF, FC, IP**Reactivity** : Human**Host species** : Rabbit**Background**

The protein encoded by this gene belongs to the highly conserved cyclin family, whose members are characterized by a dramatic periodicity in protein abundance through the cell cycle. Cyclins function as regulators of CDK kinases. Different cyclins exhibit distinct expression and degradation patterns which contribute to the temporal coordination of each mitotic event. This cyclin forms a complex with CDK4 or CDK6 and functions as a regulatory subunit of the complex, whose activity is required for cell cycle G1/S transition. This protein has been shown to interact with and be involved in the phosphorylation of tumor suppressor protein Rb. Knockout studies of the homologous gene in mouse suggest the essential roles of this gene in ovarian granulosa and germ cell proliferation. High level expression of this gene was observed in ovarian and testicular tumors. Mutations in this gene are associated with megalencephaly-polymicrogyria-polydactyly-hydrocephalus syndrome 3 (MPPH3). [provided by RefSeq, Sep 2014]

Immunogen

Recombinant protein of human Cyclin D2

Gene ID

894

Swiss Prot

P30279

Synonyms

MPPH3; KIAK0002

Reactivity

Human

Application

WB, ICC/IF, FC, IP

Recommended dilution

WB: 1:1000

ICC/IF: 1:100

FC: 1:200-1:500

IP: 1:20-1:50


Calculated MW

33 kDa

Observed MW

33 kDa

Host species	Rabbit
Clonality	Monoclonal
Clonality No.	DGR12228
Isotype	IgG
Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.



Western blot analysis of extracts from U-2OS cells using db15797 at 1:1000.