

Recombinant

DGRmAb®

Dnmt1 (DGR11950) Rabbit mAb

db15873

Package : 10µL 20µL 50µL 100µL

Product Name : Dnmt1 (DGR11950) Rabbit mAb**Cat.No.:** db15873**Synonyms** : AIM; DNMT; MCMT; CXXC9; HSN1E; ADCADN; m.HsaI**Application** : WB, IHC, ICC/IF, FC**Reactivity** : Human, Mouse, Rat**Host species** : Rabbit**Background**

This gene encodes an enzyme that transfers methyl groups to cytosine nucleotides of genomic DNA. This protein is the major enzyme responsible for maintaining methylation patterns following DNA replication and shows a preference for hemi-methylated DNA. Methylation of DNA is an important component of mammalian epigenetic gene regulation. Aberrant methylation patterns are found in human tumors and associated with developmental abnormalities. Variation in this gene has been associated with cerebellar ataxia, deafness, and narcolepsy, and neuropathy, hereditary sensory, type IE. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jan 2016]

Immunogen

A synthetic peptide of human Dnmt1

Gene ID

1786

Swiss Prot

P26358

Synonyms

AIM; DNMT; MCMT; CXXC9; HSN1E; ADCADN; m.HsaI

Reactivity

Human, Mouse, Rat

Application

WB, IHC, ICC/IF, FC

Recommended dilution

WB: 1:1000

IHC: 1:100

ICC/IF: 1:200-1:1000

FC: 1:100

Calculated MW

183 kDa

Observed MW

183 kDa


Host species

Rabbit

Clonality

Monoclonal

Clonality No.	DGR11950
Isotype	IgG
Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.



Immunofluorescence analysis of HeLa cells labelling Dnmt1 with db15873.

The cells were fixed with 4% PFA (10min, RT) followed by treatment with 0.1% Triton X-100 (10min, RT), and blocked in 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween 20 for 1h. The cells were then incubate with db15873 (1:1000) at room temprature for 1h, followed by a further incubation at room temperature for 45min with Goat Anti Rabbit IgG (H+L)-AF488 (db10005, shown in green). Nuclear DNA was labeled in blue with DAPI.

Control: Secondary antibody only.