

Recombinant

DGRmAb®

## AKT1/2 (DGR11878) Rabbit mAb

db15890

Package : 10μL 20μL 50μL 100μL

**Product Name :** AKT1/2 (DGR11878) Rabbit mAb**Cat.No.:** db15890**Synonyms :** AKT; PKB; RAC; CWS6; PRKBA; PKB-ALPHA; RAC-ALPHA**Application :** WB, IHC-P, ICC/IF, FC**Reactivity :** Human,Mouse,Rat**Host species :** Rabbit**Background**

The serine-threonine protein kinase encoded by the AKT1 gene is catalytically inactive in serum-starved primary and immortalized fibroblasts. AKT1 and the related AKT2 are activated by platelet-derived growth factor. The activation is rapid and specific, and it is abrogated by mutations in the pleckstrin homology domain of AKT1. It was shown that the activation occurs through phosphatidylinositol 3-kinase. In the developing nervous system AKT is a critical mediator of growth factor-induced neuronal survival. Survival factors can suppress apoptosis in a transcription-independent manner by activating the serine/threonine kinase AKT1, which then phosphorylates and inactivates components of the apoptotic machinery. Mutations in this gene have been associated with the Proteus syndrome. Multiple alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Jul 2011]

**Immunogen**

A synthetic peptide of human AKT1

**Gene ID**

207

**Swiss Prot**

P31749

**Synonyms**

AKT; PKB; RAC; CWS6; PRKBA; PKB-ALPHA; RAC-ALPHA

**Reactivity**

Human,Mouse,Rat

**Application**

WB, IHC-P, ICC/IF, FC

**Recommended dilution**WB: 1:1000  
IHC-P: 1:100  
ICC/IF: 1:200-1:500  
FC: 1:100-1:200**Calculated MW**

56 kDa

**Observed MW**

56 kDa

**Host species**

Rabbit

Clonality	Monoclonal
Clonality No.	DGR11878
Isotype	IgG
Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.