

Recombinant

DGRmAb®

TAB1 (DGR11733) Rabbit mAb

db15922

Package : 10μL 20μL 50μL 100μL

Product Name : TAB1 (DGR11733) Rabbit mAb**Cat.No.:** db15922**Synonyms :** 3'-Tab1; MAP3K7IP1**Application :** WB, IHC-P, ICC/IF, FC**Reactivity :** Human,Mouse,Rat**Host species :** Rabbit**Background**

The protein encoded by this gene was identified as a regulator of the MAP kinase kinase kinase MAP3K7/TAK1, which is known to mediate various intracellular signaling pathways, such as those induced by TGF beta, interleukin 1, and WNT-1. This protein interacts and thus activates TAK1 kinase. It has been shown that the C-terminal portion of this protein is sufficient for binding and activation of TAK1, while a portion of the N-terminus acts as a dominant-negative inhibitor of TGF beta, suggesting that this protein may function as a mediator between TGF beta receptors and TAK1. This protein can also interact with and activate the mitogen-activated protein kinase 14 (MAPK14/p38alpha), and thus represents an alternative activation pathway, in addition to the MAPKK pathways, which contributes to the biological responses of MAPK14 to various stimuli. Alternatively spliced transcript variants encoding distinct isoforms have been reported. [provided by RefSeq, Jul 2008]

Immunogen

A synthetic peptide of human TAB1

Gene ID

10454

Swiss Prot

Q15750

Synonyms

3'-Tab1; MAP3K7IP1

Reactivity

Human,Mouse,Rat

Application

WB, IHC-P, ICC/IF, FC


Recommended dilutionWB: 1:1000
IHC-P: 1:100-1:200
ICC/IF: 1:100-1:200
FC: 1:20**Calculated MW**

55 kDa


Observed MW

55 kDa

Host species	Rabbit
Clonality	Monoclonal
Clonality No.	DGR11733
Isotype	IgG
Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.



Western blot analysis of extracts from HeLa cells using db15922 at 1:1000.



Immunofluorescence analysis of HeLa cells labelling TAB1 with db15922.

The cells were fixed with cold 100% methanol (10min, 4℃) and blocked in 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween 20 for 1h. The cells were then incubate with db15922 (1:200) at room temprature for 1h, followed by a further incubation at room temperature for 45min with Goat Anti Rabbit IgG (H+L)-AF488 ([db10005](#), shown in green). Nuclear DNA was labeled in blue with DAPI.

Control: Secondary antibody only.