

Recombinant**DGRmAb®****ATF-4 (DGR11603) Rabbit mAb****db15953****Package : 10µL 20µL 50µL 100µL****Product Name :** ATF-4 (DGR11603) Rabbit mAb**Cat.No.:** db15953**Synonyms :** CREB2; TXREB; CREB-2; TAXREB67**Application :** WB, IHC-P, ICC/IF, FC, IP**Reactivity :** Human**Host species :** Rabbit**Background**

This gene encodes a transcription factor that was originally identified as a widely expressed mammalian DNA binding protein that could bind a tax-responsive enhancer element in the LTR of HTLV-1. The encoded protein was also isolated and characterized as the cAMP-response element binding protein 2 (CREB-2). The protein encoded by this gene belongs to a family of DNA-binding proteins that includes the AP-1 family of transcription factors, cAMP-response element binding proteins (CREBs) and CREB-like proteins. These transcription factors share a leucine zipper region that is involved in protein-protein interactions, located C-terminal to a stretch of basic amino acids that functions as a DNA binding domain. Two alternative transcripts encoding the same protein have been described. Two pseudogenes are located on the X chromosome at q28 in a region containing a large inverted duplication. [provided by RefSeq, Sep 2011]

Immunogen

Recombinant protein of human ATF-4

Gene ID

468

Swiss Prot

P18848

Synonyms

CREB2; TXREB; CREB-2; TAXREB67

Reactivity

Human

Application

WB, IHC-P, ICC/IF, FC, IP

Recommended dilution

WB: 1:1000

IHC-P: 1:50

ICC/IF: 1:200-1:1000

FC: 1:50-1:100

IP: 1:50


Calculated MW

39 kDa

Observed MW

49 kDa

| | |
|--------------------------|---|
| Host species | Rabbit |
| Clonality | Monoclonal |
| Clonality No. | DGR11603 |
| Isotype | IgG |
| Purity | Affinity Purification |
| Conjugation | Un-conjugated |
| Storage Stability | Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt. |



Immunofluorescence analysis of HeLa cells labelling ATF-4 with db15953.

The cells were fixed with 4% PFA (10min, RT) followed by treatment with 0.1% Triton X-100 (10min, RT), and blocked in 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween 20 for 1h. The cells were then incubate with db15953(1:1000) at room temprature for 1h, followed by a further incubation at room temperature for 45min with Goat Anti Rabbit IgG (H+L)-AF488 (db10005, shown in green). Nuclear DNA was labeled in blue with DAPI.

Control: Secondary antibody only.