

Recombinant

DGRmAb®

NFkB p100 (DGR11547) Rabbit mAb

db15969

Package : 10µL 20µL 50µL 100µL

Product Name : NFkB p100 (DGR11547) Rabbit mAb**Cat.No.:** db15969**Synonyms** : p52; p100; H2TF1; LYT10; CVID10; LYT-10; NF-kB2; p49/p100**Application** : WB, ICC/IF, FC, IP**Reactivity** : Human,Mouse,Rat**Host species** : Rabbit**Background**

This gene encodes a subunit of the transcription factor complex nuclear factor-kappa-B (NFkB). The NFkB complex is expressed in numerous cell types and functions as a central activator of genes involved in inflammation and immune function. The protein encoded by this gene can function as both a transcriptional activator or repressor depending on its dimerization partner. The p100 full-length protein is co-translationally processed into a p52 active form. Chromosomal rearrangements and translocations of this locus have been observed in B cell lymphomas, some of which may result in the formation of fusion proteins. There is a pseudogene for this gene on chromosome 18. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Dec 2013]

Immunogen

Recombinant protein of human NFkB p100

Gene ID

4791

Swiss Prot

Q00653

Synonyms

p52; p100; H2TF1; LYT10; CVID10; LYT-10; NF-kB2; p49/p100

Reactivity

Human,Mouse,Rat

Application

WB, ICC/IF, FC, IP

Recommended dilutionWB: 1:1000
ICC/IF: 1:100-1:200
FC: 1:50-1:100
IP: 1:20**Calculated MW**

97 kDa

Observed MW



120 kDa

Host species

Rabbit

Clonality	Monoclonal
Clonality No.	DGR11547
Isotype	IgG
Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.

Western blot analysis of extracts from HeLa, MCF-7 cells using db15969 at 1:1000.



Immunofluorescence analysis of NIH/3T3 cells labelling NFkB p100 with db15969.

The cells were fixed with 4% PFA (10min, RT) followed by treatment with 0.1% Triton X-100 (10min, RT), and blocked in 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween 20 for 1h. The cells were then incubate with db15969(1:200) at room temprature for 1h, followed by a further incubation at room temperature for 45min with Goat Anti Rabbit IgG (H+L)-AF488 (db10005, shown in green). Nuclear DNA was labeled in blue with DAPI.

Control: Secondary antibody only.