



Recombinant

DGRmAb[®]

RUNX2 (DGR11349) Rabbit mAb

db16015 Package : 10μL 20μL 50μL 100μL

Product Name: RUNX2 (DGR11349) Rabbit mAb

Cat.No.: db16015

Synonyms: CCD; AML3; CCD1; CLCD; OSF2; CBFA1; OSF-2; PEA2aA; PEBP2aA; CBF-alpha-1

Application : IHC-P, ICC/IF, FC **Reactivity :** Human, Mouse, Rat

Host species: Rabbit

Background This gene is a member of the RUNX family of transcription factors and encodes a nuclear protein

with an Runt DNA-binding domain. This protein is essential for osteoblastic differentiation and skeletal morphogenesis and acts as a scaffold for nucleic acids and regulatory factors involved in skeletal gene expression. The protein can bind DNA both as a monomer or, with more affinity, as a subunit of a heterodimeric complex. Two regions of potential trinucleotide repeat expansions are present in the N-terminal region of the encoded protein, and these and other mutations in this gene

have been associated with the bone development disorder cleidocranial dysplasia (CCD).

Transcript variants that encode different protein isoforms result from the use of alternate promoters

as well as alternate splicing. [provided by RefSeq, Jul 2016]

Immunogen Recombinant protein of human RUNX2

Gene ID 860

Swiss Prot Q13950

Synonyms CCD; AML3; CCD1; CLCD; OSF2; CBFA1; OSF-2; PEA2aA; PEBP2aA; CBF-alpha-1

Reactivity Human, Mouse, Rat

Application IHC-P, ICC/IF, FC

Recommended dilution HC-P: 1:200-1:1000

ICC/IF: 1:200-1:1000

FC: 1:50-1:100

Calculated MW 57 kDa

Host species Rabbit

Clonality Monoclonal

Clonality No. DGR11349



For Research Use Only **Product Datasheet**


Isotype IgG

Purity Affinity Purification

Conjugation Un-conjugated

Storage Stability Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium

azide and 0.05% BSA. Stable for 12 months from date of receipt.



Immunofluorescence analysis of HeLa cells labelling RUNX2 with db16015.

The cells were fixed with 4% PFA (10min, RT) followed by treatment with 0.1% Triton X-100 (10min, RT), and blocked in 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween 20 for 1h. The cells were then incubate with db16015(1:200) at room temprature for 1h, followed by a further incubation at room temperature for 45min with Goat Anti Rabbit lgG (H+L)-AF488 (db10005, shown in green). Nuclear DNA was labeled in blue with DAPI.

Control: Secondary antibody only.