

Recombinant

DGRmAb®

RUNX2 (DGR11349) Rabbit mAb

db16015

Package : 10µL 20µL 50µL 100µL

Product Name : RUNX2 (DGR11349) Rabbit mAb

Cat.No.: db16015

Synonyms : CCD; AML3; CCD1; CLCD; OSF2; CBFA1; OSF-2; PEA2aA; PEBP2aA; CBF-alpha-1

Application : IHC-P, ICC/IF, FC

Reactivity : Human,Mouse,Rat

Host species : Rabbit

Background

This gene is a member of the RUNX family of transcription factors and encodes a nuclear protein with an Runt DNA-binding domain. This protein is essential for osteoblastic differentiation and skeletal morphogenesis and acts as a scaffold for nucleic acids and regulatory factors involved in skeletal gene expression. The protein can bind DNA both as a monomer or, with more affinity, as a subunit of a heterodimeric complex. Two regions of potential trinucleotide repeat expansions are present in the N-terminal region of the encoded protein, and these and other mutations in this gene have been associated with the bone development disorder cleidocranial dysplasia (CCD). Transcript variants that encode different protein isoforms result from the use of alternate promoters as well as alternate splicing. [provided by RefSeq, Jul 2016]

Immunogen

Recombinant protein of human RUNX2

Gene ID

860

Swiss Prot

Q13950

Synonyms

CCD; AML3; CCD1; CLCD; OSF2; CBFA1; OSF-2; PEA2aA; PEBP2aA; CBF-alpha-1

Reactivity

Human,Mouse,Rat

Application

IHC-P, ICC/IF, FC

Recommended dilution

WB: 1:1000
 IHC-P: 1:200-1:1000
 ICC/IF: 1:200-1:1000
 FC: 1:50-1:100

Calculated MW

57 kDa


Observed MW

57 kDa


Host species

Rabbit

Clonality	Monoclonal
Clonality No.	DGR11349
Isotype	IgG
Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.



Western blot analysis of extracts from 3T3 cells using db16015 at 1:1000.



Immunofluorescence analysis of HeLa cells labelling RUNX2 with db16015.

The cells were fixed with 4% PFA (10min, RT) followed by treatment with 0.1% Triton X-100 (10min, RT), and blocked in 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween 20 for 1h. The cells were then incubate with db16015(1:200) at room temprature for 1h, followed by a further incubation at room temperature for 45min with Goat Anti Rabbit IgG (H+L)-AF488 (db10005, shown in green). Nuclear DNA was labeled in blue with DAPI.

Control: Secondary antibody only.