

Recombinant

DGRmAb®

ALIX (DGR35650) Rabbit mAb

db16124

Package : 10µL 20µL 50µL 100µL

Product Name : ALIX (DGR35650) Rabbit mAb**Cat.No.:** db16124**Synonyms** : AIP1; ALIX; HP95; DRIP4; MCPH29**Application** : WB, ICC/IF, FC, IP**Reactivity** : Human,Mouse,Rat**Host species** : Rabbit**Background**

This gene encodes a protein that functions within the ESCRT pathway in the abscission stage of cytokinesis, in intraluminal endosomal vesicle formation, and in enveloped virus budding. Studies using mouse cells have shown that overexpression of this protein can block apoptosis. In addition, the product of this gene binds to the product of the PDCD6 gene, a protein required for apoptosis, in a calcium-dependent manner. This gene product also binds to endophilins, proteins that regulate membrane shape during endocytosis. Overexpression of this gene product and endophilins results in cytoplasmic vacuolization, which may be partly responsible for the protection against cell death. Several alternatively spliced transcript variants encoding different isoforms have been found for this gene. Related pseudogenes have been identified on chromosome 15. [provided by RefSeq, Jan 2012]

Immunogen

Recombinant protein of human ALIX

Gene ID

10015, 18571, 501083

Swiss Prot

Q8WUM4, Q9WU78, Q9QZA2

Synonyms

AIP1; ALIX; HP95; DRIP4; MCPH29

Reactivity

Human,Mouse,Rat

Application

WB, ICC/IF, FC, IP

Recommended dilution

WB: 1:1000

ICC/IF: 1:500-1:5000

FC: 1:200-1:500

IP: 1:20-1:50

Calculated MW

96 kDa


Observed MW

90-100 kDa

Host species

Rabbit


Clonality	Monoclonal
Clonality No.	DGR35650
Isotype	IgG
Purity	Affinity Purification
Conjugation	Un-conjugated
Storage Stability	Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.



Immunofluorescence analysis of HeLa cells labelling ALIX with db16124.

The cells were fixed with cold 100% methanol (10min, 4℃) and blocked in 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween 20 for 1h. The cells were then incubate with db16124 (1:500) at room temprature for 1h, followed by a further incubation at room temperature for 45min with Goat Anti Rabbit IgG (H+L)-AF488 (db10005, shown in green). Nuclear DNA was labeled in blue with DAPI.

Control: Secondary antibody only.



Western blot analysis of extracts from HeLa cells using db16124 at 1:1000.