

Recombinant

DGRmAb®

ACAA2 (DGR36015) Rabbit mAb

db16125


Package : 10µL 20µL 50µL 100µL

Product Name : ACAA2 (DGR36015) Rabbit mAb**Cat.No.:** db16125**Synonyms** : T1; DSAEC**Application** : WB, IHC-P, ICC/IF**Reactivity** : Human,Mouse,Rat**Host species** : Rabbit


Background	The encoded protein catalyzes the last step of the mitochondrial fatty acid beta-oxidation spiral. Unlike most mitochondrial matrix proteins, it contains a non-cleavable amino-terminal targeting signal. [provided by RefSeq, Jul 2008]
Immunogen	A synthetic peptide of human ACAA2
Gene ID	10449, 52538, 170465
Swiss Prot	P42765, Q8BWT1, P13437
Synonyms	T1; DSAEC
Reactivity	Human,Mouse,Rat
Application	WB, IHC-P, ICC/IF
Recommended dilution	WB: 1:1000-1:5000 IHC-P: 1:100-1:200 ICC/IF: 1:100-1:200
Calculated MW	42 kDa
Observed MW	42 kDa
Host species	Rabbit
Clonality	Monoclonal
Clonality No.	DGR36015
Isotype	IgG
Purity	Affinity Purification
Conjugation	Un-conjugated

Storage Stability

Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.



Western blot analysis of extracts from HepG2 cells using db16125 at 1:1000.



Immunofluorescence analysis of HepG2 cells labelling ACAA2 with db16125.

The cells were fixed with 4% PFA (10min, RT) followed by treatment with 0.1% Triton X-100 (10min, RT), and blocked in 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween 20 for 1h. The cells were then incubate with db16125 (1:100) at room temprature for 1h, followed by a further incubation at room temperature for 45min with Goat Anti Rabbit IgG (H+L)-AF488 (db10005, shown in green). Nuclear DNA was labeled in blue with DAPI.

Control: Secondary antibody only.