



## Recombinant



## Acetyl-Histone H3 (Lys27) (DGR14746) Rabbit mAb (PBS Only)

db11759-PBS Package : 100μg

Product Name: Acetyl-Histone H3 (Lys27) (DGR14746) Rabbit mAb (PBS Only)

Cat.No.: db11759-PBS

Synonyms: H3/j; H3C1; H3C2; H3C3; H3C4; H3C6; H3C7; H3C8; H3FJ; H3C10; H3C11; HIST1H3J

**Application**: WB, IHC-P, ICC/IF, FC **Reactivity**: Human, Mouse, Rat

Host species: Rabbit

**Background** Histones are basic nuclear proteins that are responsible for the nucleosome structure of the

chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene is intronless

and encodes a replication-dependent histone that is a member of the histone H3 family.

Transcripts from this gene lack polyA tails but instead contain a palindromic termination element. This gene is found in the small histone gene cluster on chromosome 6p22-p21.3. [provided by

RefSeq, Aug 2015]

**Immunogen** A synthetic acetyl-peptide corresponding to residues surrounding Lys27 of human Histone H3

Gene ID 8356

Swiss Prot P68431

**Synonyms** H3/j; H3C1; H3C2; H3C3; H3C4; H3C6; H3C7; H3C8; H3FJ; H3C10; H3C11; HIST1H3J

Reactivity Human.Mouse.Rat

**Application** WB, IHC-P, ICC/IF, FC

Recommended dilution WB: 1:5000-1:50000

IHC-P: 1:1000-1:10000 ICC/IF: 1:500-1: 5000

FC: 1:200-1:1000

Calculated MW 15 kDa

Observed MW 17 kDa

Host species Rabbit



## For Research Use Only **Product Datasheet**

**Clonality** Monoclonal

Clonality No. DGR14746

**Isotype** IgG

**Purity** Affinity Purification


**Conjugation** Un-conjugated

Concentration 1 mg/ml


Formulation PBS Only

Storage Stability Store at -20°C. Recommended to aliquot into single-use vials. Supplied in 1X PBS (pH 7.4). BSA

and Azide Free. Stable for 12 months from date of receipt.



Western blot detection of Histone H3 (acetyl-Lys27) in Hela, A549, HL-60 cell lysates using Histone H3 (acetyl-Lys27) antibody(1:1000 diluted).



10 -

Western blot analysis of extracts from C6 cells using db11759