

Recombinant

DGRmAb®

Proteasome 20S LMP2 (DGR18898) Rabbit mAb (PBS Only)

db14614-PBS

Package : 10µg 100µg

Product Name : Proteasome 20S LMP2 (DGR18898) Rabbit mAb (PBS Only)**Cat.No.:** db14614-PBS**Synonyms** : LMP2; PSMB6i; RING12; beta1i**Application** : WB, ICC/IF, FC, IP**Reactivity** : Human**Host species** : Rabbit**Background**

The proteasome is a multicatalytic proteinase complex with a highly ordered ring-shaped 20S core structure. The core structure is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. This gene encodes a member of the proteasome B-type family, also known as the T1B family, that is a 20S core beta subunit. This gene is located in the class II region of the MHC (major histocompatibility complex). Expression of this gene is induced by gamma interferon and this gene product replaces catalytic subunit 1 (proteasome beta 6 subunit) in the immunoproteasome. Proteolytic processing is required to generate a mature subunit. [provided by RefSeq, Mar 2010]

Immunogen

A synthetic peptide of human Proteasome 20S LMP2

Gene ID

5698

Swiss Prot

P28065

Synonyms

LMP2; PSMB6i; RING12; beta1i

Reactivity

Human

Application

WB, ICC/IF, FC, IP

Calculated MW

23 kDa

Observed MW

23 kDa

Host species

Rabbit

Clonality

Monoclonal

Clonality No.

DGR18898

Isotype	IgG
Purity	Affinity Purification
Conjugation	Un-conjugated
Concentration	1 mg/mL
Formulation	PBS Only
Storage Stability	Store at -20°C. Recommended to aliquot into single-use vials. Supplied in 1X PBS (pH 7.4). BSA and Azide Free. Stable for 12 months from date of receipt.