

Recombinant

DGRmAb®

Phospho-gamma H2A.X (Ser139) (DGR18728) Rabbit mAb (PBS Only)

db13191-PBS

Package : 10µg 100µg

Product Name : Phospho-gamma H2A.X (Ser139) (DGR18728) Rabbit mAb (PBS Only)**Cat.No.:** db13191-PBS**Synonyms** : H2AX; H2A.X; H2A/X**Application** : WB, IHC-P, ICC/IF, IP**Reactivity** : Human,Mouse,Rat**Host species** : Rabbit**Background**

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene encodes a replication-independent histone that is a member of the histone H2A family, and generates two transcripts through the use of the conserved stem-loop termination motif, and the polyA addition motif. [provided by RefSeq, Oct 2015]

Immunogen

A synthetic phosphopeptide corresponding to residues surrounding Ser139 of human gamma H2A.X

Gene ID

3014

Swiss Prot

P16104

Synonyms

H2AX; H2A.X; H2A/X

Reactivity

Human,Mouse,Rat

Application

WB, IHC-P, ICC/IF, IP

Calculated MW

15 kDa

Observed MW

15 kDa

Host species

Rabbit

Clonality

Monoclonal


Clonality No.

DGR18728


Isotype

IgG

Purity	Affinity Purification
Conjugation	Un-conjugated
Concentration	1 mg/mL
Formulation	PBS Only
Storage Stability	Store at -20°C. Recommended to aliquot into single-use vials. Supplied in 1X PBS (pH 7.4). BSA and Azide Free. Stable for 12 months from date of receipt.



Western blot analysis of extracts from Jurkat, THP-1, C6, 3T3 cells using [db13191](#) at 1:1000.



Western blot analysis of extracts from HepG2 cells using [db13191](#) at 1:1000.