



Recombinant

DGRmAb[®]

PI3-Kinase p85 alpha (DGR16380) Rabbit mAb (PBS Only)

db16082-PBS Package : 100μg

Product Name: PI3-Kinase p85 alpha (DGR16380) Rabbit mAb (PBS Only)

Cat.No.: db16082-PBS

Synonyms: p85; AGM7; GRB1; IMD36; p85-ALPHA

Application : IHC, ICC/IF, FC **Reactivity :** Human, Mouse, Rat

Host species: Rabbit

Background Phosphatidylinositol 3-kinase phosphorylates the inositol ring of phosphatidylinositol at the 3-prime

position. The enzyme comprises a 110 kD catalytic subunit and a regulatory subunit of either 85, 55, or 50 kD. This gene encodes the 85 kD regulatory subunit. Phosphatidylinositol 3-kinase plays

an important role in the metabolic actions of insulin, and a mutation in this gene has been

associated with insulin resistance. Alternative splicing of this gene results in four transcript variants

encoding different isoforms. [provided by RefSeq, Jun 2011]

Immunogen A synthetic peptide of human PI3 Kinase p85 alpha

Gene ID 5295

Swiss Prot P27986

Synonyms p85; AGM7; GRB1; IMD36; p85-ALPHA

Reactivity Human, Mouse, Rat

Application IHC, ICC/IF, FC

Recommended dilution IHC: 1:100

ICC/IF: 1:50

FC: 1:50

Calculated MW 84 kDa

Observed MW 85 kDa

Host species Rabbit

Clonality Monoclonal

Clonality No. DGR16380

Isotype IgG



For Research Use Only **Product Datasheet**

Purity Affinity Purification


Conjugation Un-conjugated

Concentration 1 mg/ml

Formulation PBS Only

Storage Stability Store at -20°C. Recommended to aliquot into single-use vials. Supplied in 1X PBS (pH 7.4). BSA

and Azide Free. Stable for 12 months from date of receipt.



Immunofluorescence analysis of 3T3 cells labelling Pl3-Kinase p85 alpha with db16082.

The cells were fixed with cold 100% methanol (10min, 4°C) and blocked in 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween 20 for 1h. The cells were then incubate with db16082 (1:50) at room temperature for 1h, followed by a further incubation at room temperature for 45min with Goat Anti Rabbit IgG (H+L)-AF488 (db10005, shown in green). Nuclear DNA was labeled in blue with DAPI.

Control: Secondary antibody only.