

Recombinant

DGRmAb®

## p53 (DGR13917) Rabbit mAb (PBS Only)

db11669-PBS

Package : 100µg

**Product Name** : p53 (DGR13917) Rabbit mAb (PBS Only)**Cat.No.:** db11669-PBS**Synonyms** : P53; BCC7; LFS1; TRP53**Application** : WB, IHC-P, ICC/IF, FC, IP**Reactivity** : Human**Host species** : Rabbit**Background**

This gene encodes tumor protein p53, which responds to diverse cellular stresses to regulate target genes that induce cell cycle arrest, apoptosis, senescence, DNA repair, or changes in metabolism. p53 protein is expressed at low level in normal cells and at a high level in a variety of transformed cell lines, where it's believed to contribute to transformation and malignancy. p53 is a DNA-binding protein containing transcription activation, DNA-binding, and oligomerization domains. It is postulated to bind to a p53-binding site and activate expression of downstream genes that inhibit growth and/or invasion, and thus function as a tumor suppressor. Mice deficient for this gene are developmentally normal but are susceptible to spontaneous tumors. Evidence to date shows that this gene contains one promoter, in contrast to alternative promoters of the human gene, and transcribes a few of splice variants which encode different isoforms, although the biological validity or the full-length nature of some variants has not been determined. [provided by RefSeq, Jul 2008]

**Immunogen**

A synthetic peptide of human p53

**Gene ID**

7157

**Swiss Prot**

P04637

**Synonyms**

P53; BCC7; LFS1; TRP53

**Reactivity**

Human

**Application**

WB, IHC-P, ICC/IF, FC, IP

**Calculated MW**

44 kDa

**Observed MW**

53 kDa


**Host species**

Rabbit


**Clonality**

Monoclonal

<b>Clonality No.</b>	DGR13917
<b>Isotype</b>	IgG
<b>Purity</b>	Affinity Purification
<b>Conjugation</b>	Un-conjugated
<b>Concentration</b>	1 mg/mL
<b>Formulation</b>	PBS Only
<b>Storage Stability</b>	Store at -20°C. Recommended to aliquot into single-use vials. Supplied in 1X PBS (pH 7.4). BSA and Azide Free. Stable for 12 months from date of receipt.



Western blot analysis of extracts from A431 cells using [db11669](#) at 1:1000.



Western blot analysis of extracts from Raji cells using [db11669](#) at 1:1000.