Recombinant DGRmAb®


Histone H3 (DGR12083) Rabbit mAb (PBS Only)

db12105-PBS **Package : 10μg 100μg**


Product Name : Histone H3 (DGR12083) Rabbit mAb (PBS Only)
Cat.No.: db12105-PBS
Synonyms : H3/A; H3FA
Application : WB, IHC-P, ICC/IF, FC
Reactivity : Human,Mouse,Rat
Host species : Rabbit

Background	Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. This structure consists of approximately 146 bp of DNA wrapped around a nucleosome, an octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H3 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element. This gene is found in the large histone gene cluster on chromosome 6p22-p21.3. [provided by RefSeq, Aug 2015]
Immunogen	A synthetic peptide of human Histone H3
Gene ID	8350
Swiss Prot	P68431
Synonyms	H3/A; H3FA
Reactivity	Human,Mouse,Rat
Application	WB, IHC-P, ICC/IF, FC
Calculated MW	15 kDa
Observed MW	17 kDa
Host species	Rabbit
Clonality	Monoclonal
Clonality No.	DGR12083
Isotype	IgG


Purity	Affinity Purification
Conjugation	Un-conjugated
Concentration	1 mg/mL
Formulation	PBS Only
Storage Stability	Store at -20°C. Recommended to aliquot into single-use vials. Supplied in 1X PBS (pH 7.4). BSA and Azide Free. Stable for 12 months from date of receipt.




Western blot analysis of extracts from Jurkat, THP-1, C6, 3T3 cells using db12105 at 1:1000.



Western blot analysis of extracts from Raji cells using db12105 at 1:1000.



Immunohistochemical analysis of paraffin-embedded human lung cancer using db12105 antibody.



Immunofluorescence analysis of HeLa cells labelling Histone H3 with db12105.

The cells were fixed with cold 100% methanol (10min, 4℃) and blocked in 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween 20 for 1h. The cells were then incubate with db12105 (1:200) at room temprature for 1h, followed by a further incubation at room temperature for 45min with Goat Anti Rabbit IgG (H+L)-AF488 (db10005, shown in green). Nuclear DNA was labeled in blue with DAPI.

Control: Secondary antibody only.