

Recombinant

DGRmAb®

C7 (DGR11341) Rabbit mAb (PBS Only)

db16016-PBS

Package : 10µg 100µg

Product Name : C7 (DGR11341) Rabbit mAb (PBS Only)**Cat.No.:** db16016-PBS**Synonyms** : Complement component C7**Application** : WB, IHC-P, ICC/IF**Reactivity** : Human,Mouse,Rat**Host species** : Rabbit**Background**

This gene encodes a serum glycoprotein that forms a membrane attack complex together with complement components C5b, C6, C8, and C9 as part of the terminal complement pathway of the innate immune system. The protein encoded by this gene contains a cholesterol-dependent cytolysin/membrane attack complex/perforin-like (CDC/MACPF) domain and belongs to a large family of structurally related molecules that form pores involved in host immunity and bacterial pathogenesis. This protein initiates membrane attack complex formation by binding the C5b-C6 subcomplex and inserts into the phospholipid bilayer, serving as a membrane anchor. Mutations in this gene are associated with a rare disorder called C7 deficiency. [provided by RefSeq, Nov 2016]

Immunogen

A synthetic peptide of human C7

Gene ID

730

Swiss Prot

P10643

Synonyms

Complement component C7

Reactivity

Human,Mouse,Rat

Application

WB, IHC-P, ICC/IF

Calculated MW

94 kDa

Observed MW

94 kDa

Host species

Rabbit

Clonality

Monoclonal


Clonality No.

DGR11341

Isotype

IgG


Purity	Affinity Purification
Conjugation	Un-conjugated
Concentration	1 mg/mL
Formulation	PBS Only
Storage Stability	Store at -20°C. Recommended to aliquot into single-use vials. Supplied in 1X PBS (pH 7.4). BSA and Azide Free. Stable for 12 months from date of receipt.



Immunofluorescence analysis of NIH/3T3 cells labelling C7 with **db16016**.

The cells were fixed with cold 100% methanol (10min, 4°C) and blocked in 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween 20 for 1h. The cells were then incubate with **db16016** (1:200) at room temperature for 1h, followed by a further incubation at room temperature for 45min with Goat Anti Rabbit IgG (H+L)-AF488 **db10005**, shown in green). Nuclear DNA was labeled in blue with DAPI.

Control: Secondary antibody only.



Western blot analysis of extracts from HepG2 cells using **db16016** at 1:1000.