

Recombinant**DGRmAb®****ICAD (DGR13646) Rabbit mAb (PBS Only)****db13371-PBS****Package : 100µg****Product Name :** ICAD (DGR13646) Rabbit mAb (PBS Only)**Cat.No.:** db13371-PBS**Synonyms :** DFF1; ICAD; DFF-45**Application :** WB**Reactivity :** Human,Mouse,Rat**Host species :** Rabbit**Background**

Apoptosis is a cell death process that removes toxic and/or useless cells during mammalian development. The apoptotic process is accompanied by shrinkage and fragmentation of the cells and nuclei and degradation of the chromosomal DNA into nucleosomal units. DNA fragmentation factor (DFF) is a heterodimeric protein of 40-kD (DFFB) and 45-kD (DFFA) subunits. DFFA is the substrate for caspase-3 and triggers DNA fragmentation during apoptosis. DFF becomes activated when DFFA is cleaved by caspase-3. The cleaved fragments of DFFA dissociate from DFFB, the active component of DFF. DFFB has been found to trigger both DNA fragmentation and chromatin condensation during apoptosis. Two alternatively spliced transcript variants encoding distinct isoforms have been found for this gene. [provided by RefSeq, Jul 2008]

Immunogen

A synthetic peptide of human ICAD

Gene ID

1676

Swiss Prot

O00273

Synonyms

DFF1; ICAD; DFF-45

Reactivity

Human,Mouse,Rat

Application

WB

Calculated MW

37 kDa

Observed MW

45 kDa

Host species

Rabbit

Clonality

Monoclonal


Clonality No.

DGR13646

Isotype

IgG

Purity	Affinity Purification
Conjugation	Un-conjugated
Concentration	1 mg/mL
Formulation	PBS Only
Storage Stability	Store at -20°C. Recommended to aliquot into single-use vials. Supplied in 1X PBS (pH 7.4). BSA and Azide Free. Stable for 12 months from date of receipt.



Western blot detection of ICAD in Rat Brain, MCF7, C6 cell lysates using ICAD antibody(1:1000 diluted).