Recombinant DGRmAb®


Securin (DGR16413) Rabbit mAb (PBS Only)

db11881-PBS **Package : 10μg 100μg**


Product Name : Securin (DGR16413) Rabbit mAb (PBS Only)
Cat.No.: db11881-PBS
Synonyms : EAP1; PTTG; HPTTG; TUTR1
Application : WB, IHC-P, ICC/IF, FC
Reactivity : Human
Host species : Rabbit

Background	The encoded protein is a homolog of yeast securin proteins, which prevent separins from promoting sister chromatid separation. It is an anaphase-promoting complex (APC) substrate that associates with a separin until activation of the APC. The gene product has transforming activity in vitro and tumorigenic activity in vivo, and the gene is highly expressed in various tumors. The gene product contains 2 PXXP motifs, which are required for its transforming and tumorigenic activities, as well as for its stimulation of basic fibroblast growth factor expression. It also contains a destruction box (D box) that is required for its degradation by the APC. The acidic C-terminal region of the encoded protein can act as a transactivation domain. The gene product is mainly a cytosolic protein, although it partially localizes in the nucleus. Three transcript variants encoding the same protein have been found for this gene. [provided by RefSeq, Sep 2013]
Immunogen	A synthetic peptide of human Securin
Gene ID	9232
Swiss Prot	O95997
Synonyms	EAP1; PTTG; HPTTG; TUTR1
Reactivity	Human
Application	WB, IHC-P, ICC/IF, FC
Calculated MW	22 kDa
Observed MW	25 kDa
Host species	Rabbit
Clonality	Monoclonal
Clonality No.	DGR16413
Isotype	IgG


Purity	Affinity Purification
Conjugation	Un-conjugated
Concentration	1 mg/mL
Formulation	PBS Only
Storage Stability	Store at -20°C. Recommended to aliquot into single-use vials. Supplied in 1X PBS (pH 7.4). BSA and Azide Free. Stable for 12 months from date of receipt.



Western blot analysis of extracts from K562, 3T3, HeLa cells using db11881 at 1:1000.



Immunohistochemical analysis of paraffin-embedded human tonsil using db11881 antibody.



Immunofluorescence analysis of HeLa cells labelling Securin with db11881.

The cells were fixed with 4% PFA (10min, RT) followed by treatment with 0.1% Triton X-100 (10min, RT), and blocked in 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween 20 for 1h. The cells were then incubate with db11881 (1:50) at room temprature for 1h, followed by a further incubation at room temperature for 45min with Goat Anti Rabbit IgG (H+L)-AF488 (db10005, shown in green). Nuclear DNA was labeled in blue with DAPI.

Control: Secondary antibody only.