

Recombinant

DGRmAb®

Securin (DGR16413) Rabbit mAb (PBS Only)

db11881-PBS

Package : 10µg 100µg

Product Name : Securin (DGR16413) Rabbit mAb (PBS Only)**Cat.No.:** db11881-PBS**Synonyms** : EAP1; PTTG; HPTTG; TUTR1**Application** : WB, IHC-P, ICC/IF, FC**Reactivity** : Human**Host species** : Rabbit**Background**

The encoded protein is a homolog of yeast securin proteins, which prevent separins from promoting sister chromatid separation. It is an anaphase-promoting complex (APC) substrate that associates with a separin until activation of the APC. The gene product has transforming activity in vitro and tumorigenic activity in vivo, and the gene is highly expressed in various tumors. The gene product contains 2 PXXP motifs, which are required for its transforming and tumorigenic activities, as well as for its stimulation of basic fibroblast growth factor expression. It also contains a destruction box (D box) that is required for its degradation by the APC. The acidic C-terminal region of the encoded protein can act as a transactivation domain. The gene product is mainly a cytosolic protein, although it partially localizes in the nucleus. Three transcript variants encoding the same protein have been found for this gene. [provided by RefSeq, Sep 2013]

Immunogen

A synthetic peptide of human Securin

Gene ID

9232

Swiss Prot

O95997

Synonyms

EAP1; PTTG; HPTTG; TUTR1

Reactivity

Human

Application

WB, IHC-P, ICC/IF, FC

Calculated MW

22 kDa

Observed MW

25 kDa

Host species

Rabbit

Clonality

Monoclonal

Clonality No.


DGR16413

Isotype


IgG

Purity	Affinity Purification
Conjugation	Un-conjugated
Concentration	1 mg/mL
Formulation	PBS Only
Storage Stability	Store at -20°C. Recommended to aliquot into single-use vials. Supplied in 1X PBS (pH 7.4). BSA and Azide Free. Stable for 12 months from date of receipt.


Western blot analysis of extracts from K562, 3T3, HeLa cells using [db11881](#) at 1:1000.



Immunohistochemical analysis of paraffin-embedded human tonsil using [db11881](#) antibody.



Immunofluorescence analysis of HeLa cells labelling Securin with [db11881](#).



The cells were fixed with 4% PFA (10min, RT) followed by treatment with 0.1% Triton X-100 (10min, RT), and blocked in 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween 20 for 1h. The cells were then incubate with [db11881](#) (1:50) at room temprature for 1h, followed by a further incubation at room temperature for 45min with Goat Anti Rabbit IgG (H+L)-AF488 ([db10005](#), shown in green). Nuclear DNA was labeled in blue with DAPI.

Control: Secondary antibody only.