

Recombinant**DGRmAb®****ICAD (DGR33010) Rabbit mAb (PBS Only)****db13280-PBS****Package : 100µg****Product Name :** ICAD (DGR33010) Rabbit mAb (PBS Only)**Cat.No.:** db13280-PBS**Synonyms :** DFF1; ICAD; DFF-45**Application :** WB, IHC-P, ICC/IF**Reactivity :** Human**Host species :** Rabbit**Background**

Apoptosis is a cell death process that removes toxic and/or useless cells during mammalian development. The apoptotic process is accompanied by shrinkage and fragmentation of the cells and nuclei and degradation of the chromosomal DNA into nucleosomal units. DNA fragmentation factor (DFF) is a heterodimeric protein of 40-kD (DFFB) and 45-kD (DFFA) subunits. DFFA is the substrate for caspase-3 and triggers DNA fragmentation during apoptosis. DFF becomes activated when DFFA is cleaved by caspase-3. The cleaved fragments of DFFA dissociate from DFFB, the active component of DFF. DFFB has been found to trigger both DNA fragmentation and chromatin condensation during apoptosis. Two alternatively spliced transcript variants encoding distinct isoforms have been found for this gene. [provided by RefSeq, Jul 2008]

Immunogen

A synthetic peptide of human ICAD

Gene ID

1676

Swiss Prot

O00273

Synonyms

DFF1; ICAD; DFF-45

Reactivity

Human

Application

WB, IHC-P, ICC/IF

Calculated MW

37 kDa

Observed MW

45 kDa

Host species

Rabbit

Clonality

Monoclonal

Clonality No.

DGR33010

Isotype


IgG

Purity	Affinity Purification
Conjugation	Un-conjugated
Concentration	1 mg/mL
Formulation	PBS Only
Storage Stability	Store at -20°C. Recommended to aliquot into single-use vials. Supplied in 1X PBS (pH 7.4). BSA and Azide Free. Stable for 12 months from date of receipt.

HeLa

Western blot analysis of extracts from HeLa cells using [db13280](#) at 1:1000.

kDa
180 -
130 -
100 -
70 -
55 -
40 -
35 -
25 -
15 -
10 -

Immunofluorescence analysis of HeLa cells labelling ICAD with [db13280](#).

The cells were fixed with cold 100% methanol (10min, 4°C) and blocked in 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween 20 for 1h. The cells were then incubate with [db13280](#) (1:200) at room temprature for 1h, followed by a further incubation at room temperature for 45min with Goat Anti Rabbit IgG (H+L)-AF488 [db10005](#), shown in green). Nuclear DNA was labeled in blue with DAPI.

Control: Secondary antibody only.