

Recombinant

DGRmAb®

EWSR1 (DGR13368) Rabbit mAb (PBS Only)

db11763-PBS

Package : 10µg 100µg

Product Name : EWSR1 (DGR13368) Rabbit mAb (PBS Only)**Cat.No.:** db11763-PBS**Synonyms** : EWS; EWS-FLI1; bK984G1.4**Application** : WB, IHC-P**Reactivity** : Human,Mouse,Rat**Host species** : Rabbit**Background**

This gene encodes a multifunctional protein that is involved in various cellular processes, including gene expression, cell signaling, and RNA processing and transport. The protein includes an N-terminal transcriptional activation domain and a C-terminal RNA-binding domain. Chromosomal translocations between this gene and various genes encoding transcription factors result in the production of chimeric proteins that are involved in tumorigenesis. These chimeric proteins usually consist of the N-terminal transcriptional activation domain of this protein fused to the C-terminal DNA-binding domain of the transcription factor protein. Mutations in this gene, specifically a t(11;22)(q24;q12) translocation, are known to cause Ewing sarcoma as well as neuroectodermal and various other tumors. Alternative splicing of this gene results in multiple transcript variants. Related pseudogenes have been identified on chromosomes 1 and 14. [provided by RefSeq, Jul 2009]

Immunogen

A synthetic peptide of human EWSR1

Gene ID

2130

Swiss Prot

Q01844

Synonyms

EWS; EWS-FLI1; bK984G1.4

Reactivity

Human,Mouse,Rat

Application

WB, IHC-P

Calculated MW

69 kDa

Observed MW

85 kDa

Host species

Rabbit


Clonality

Monoclonal

Clonality No.

DGR13368

Isotype	IgG
Purity	Affinity Purification
Conjugation	Un-conjugated
Concentration	1 mg/mL
Formulation	PBS Only
Storage Stability	Store at -20°C. Recommended to aliquot into single-use vials. Supplied in 1X PBS (pH 7.4). BSA and Azide Free. Stable for 12 months from date of receipt.



Western blot detection of EWSR1 in HeLa cell lysates using EWSR1 antibody(1:1000 diluted).