

Recombinant

DGRmAb®

NF-κB p65 (DGR19469) Rabbit mAb (PBS Only)

db13524-PBS

Package : 100μg

Product Name : NF-κB p65 (DGR19469) Rabbit mAb (PBS Only)**Cat.No.:** db13524-PBS**Synonyms** : p65; NFKB3**Application** : WB, ICC/IF, FC**Reactivity** : Human, Mouse, Rat**Host species** : Rabbit**Background**

NF-kappa-B is a ubiquitous transcription factor involved in several biological processes. It is held in the cytoplasm in an inactive state by specific inhibitors. Upon degradation of the inhibitor, NF-kappa-B moves to the nucleus and activates transcription of specific genes. NF-kappa-B is composed of NFKB1 or NFKB2 bound to either REL, RELA, or RELB. The most abundant form of NF-kappa-B is NFKB1 complexed with the product of this gene, RELA. Four transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Sep 2011]

Immunogen

A synthetic peptide of human NF-κB p65

Gene ID

5970

Swiss Prot

Q04206

Synonyms

p65; NFKB3

Reactivity

Human, Mouse, Rat

Application

WB, ICC/IF, FC

Calculated MW

60 kDa

Observed MW

65 kDa

Host species

Rabbit

Clonality

Monoclonal

Clonality No.

DGR19469

Isotype

IgG


Purity

Affinity Purification


Conjugation

Un-conjugated

Concentration	1 mg/mL
Formulation	PBS Only
Storage Stability	Store at -20°C. Recommended to aliquot into single-use vials. Supplied in 1X PBS (pH 7.4). BSA and Azide Free. Stable for 12 months from date of receipt.



Western blot analysis of extracts from Jurkat, C6 cells using [db13524](#) at 1:1000.



Immunofluorescence analysis of HeLa cells labelling NF-KB p65 with [db13524](#).

The cells were fixed with cold 100% methanol (10min, 4°C) and blocked in 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween 20 for 1h. The cells were then incubate with [db13524](#) (1:100) at room temprature for 1h, followed by a further incubation at room temperature for 45min with Goat Anti Rabbit IgG (H+L)-AF488 ([db10005](#), shown in green). Nuclear DNA was labeled in blue with DAPI.

Control: Secondary antibody only.