Recombinant DGRmAb®


Cyclin H (DGR13708) Rabbit mAb (PBS Only)

db14756-PBS **Package : 10μg 100μg**

Product Name : Cyclin H (DGR13708) Rabbit mAb (PBS Only)
Cat.No.: db14756-PBS
Synonyms : CAK; p34; p37; CycH
Application : WB
Reactivity : Human,Mouse,Rat
Host species : Rabbit

Background	The protein encoded by this gene belongs to the highly conserved cyclin family, whose members are characterized by a dramatic periodicity in protein abundance through the cell cycle. Cyclins function as regulators of CDK kinases. Different cyclins exhibit distinct expression and degradation patterns which contribute to the temporal coordination of each mitotic event. This cyclin forms a complex with CDK7 kinase and ring finger protein MAT1. The kinase complex is able to phosphorylate CDK2 and CDC2 kinases, thus functions as a CDK-activating kinase (CAK). This cyclin and its kinase partner are components of TFIIH, as well as RNA polymerase II protein complexes. They participate in two different transcriptional regulation processes, suggesting an important link between basal transcription control and the cell cycle machinery. A pseudogene of this gene is found on chromosome 4. Alternate splicing results in multiple transcript variants. [provided by RefSeq, Nov 2010]
Immunogen	A synthetic peptide of human Cyclin H
Gene ID	902
Swiss Prot	P51946
Synonyms	CAK; p34; p37; CycH
Reactivity	Human,Mouse,Rat
Application	WB
Calculated MW	38 kDa
Observed MW	36 kDa
Host species	Rabbit
Clonality	Monoclonal
Clonality No.	DGR13708

Isotype	IgG
Purity	Affinity Purification
Conjugation	Un-conjugated
Concentration	1 mg/mL
Formulation	PBS Only
Storage Stability	Store at -20°C. Recommended to aliquot into single-use vials. Supplied in 1X PBS (pH 7.4). BSA and Azide Free. Stable for 12 months from date of receipt.



Western blot detection of Cyclin H in Jurkat cell lysates using Cyclin H antibody(1:1000 diluted).