

Recombinant

DGRmAb®

## p21 (DGR15110) Rabbit mAb (PBS Only)

db15140-PBS

Package : 100µg

**Product Name** : p21 (DGR15110) Rabbit mAb (PBS Only)**Cat.No.:** db15140-PBS**Synonyms** : P21; CDKI; CIP1; SDI1; Waf1; mda6; CAP20; Cdkn1; p21WAF; p21Cip1**Application** : WB, IHC-P, ICC/IF, FC, IP**Reactivity** : Mouse**Host species** : Rabbit**Background**

This gene encodes a potent cyclin-dependent kinase inhibitor. The encoded protein binds to and inhibits the activity of cyclin-cyclin-dependent kinase2 or cyclin-dependent kinase4 complexes, and thus functions as a regulator of cell cycle progression at the G1 phase. The expression of this gene is tightly controlled by the tumor suppressor protein p53, through which this protein mediates the p53-dependent cell cycle G1 phase arrest in response to a variety of stress stimuli. This protein can interact with proliferating cell nuclear antigen, a DNA polymerase accessory factor, and plays a regulatory role in S phase DNA replication and DNA damage repair. This protein was reported to be specifically cleaved by CASP3-like caspases, which thus leads to a dramatic activation of cyclin-dependent kinase2, and may be instrumental in the execution of apoptosis following caspase activation. Mice that lack this gene have the ability to regenerate damaged or missing tissue. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Sep 2015]

**Immunogen**

Recombinant protein of mouse p21

**Gene ID**

12575

**Swiss Prot**

P39689

**Synonyms**

P21; CDKI; CIP1; SDI1; Waf1; mda6; CAP20; Cdkn1; p21WAF; p21Cip1

**Reactivity**

Mouse

**Application**

WB, IHC-P, ICC/IF, FC, IP

**Calculated MW**

18 kDa

**Observed MW**

18 kDa

**Host species**

Rabbit


**Clonality**

Monoclonal

**Clonality No.**

DGR15110

<b>Isotype</b>	IgG
<b>Purity</b>	Affinity Purification
<b>Conjugation</b>	Un-conjugated
<b>Concentration</b>	1 mg/mL
<b>Formulation</b>	PBS Only
<b>Storage Stability</b>	Store at -20°C. Recommended to aliquot into single-use vials. Supplied in 1X PBS (pH 7.4). BSA and Azide Free. Stable for 12 months from date of receipt.



Immunofluorescence analysis of 3T3 cells labelling p21 with [db15140](#).

The cells were fixed with 4% PFA (10min, RT) followed by treatment with 0.1% Triton X-100 (10min, RT), and blocked in 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween 20 for 1h. The cells were then incubate with [db15140](#) (1:200) at room temperature for 1h, followed by a further incubation at room temperature for 45min with Goat Anti Rabbit IgG (H+L)-AF488 [db10005](#), shown in green). Nuclear DNA was labeled in blue with DAPI.

Control: Secondary antibody only.