

Recombinant

DGRmAb®

MEK1 (DGR11737) Rabbit mAb (PBS Only)

db15920-PBS

Package : 10µg 100µg

Product Name : MEK1 (DGR11737) Rabbit mAb (PBS Only)**Cat.No.:** db15920-PBS**Synonyms** : CFC3; MEK1; MKK1; MAPKK1; PRKMK1**Application** : WB, IHC-P, ICC/IF, FC, IP**Reactivity** : Human**Host species** : Rabbit**Background**

The protein encoded by this gene is a member of the dual specificity protein kinase family, which acts as a mitogen-activated protein (MAP) kinase kinase. MAP kinases, also known as extracellular signal-regulated kinases (ERKs), act as an integration point for multiple biochemical signals. This protein kinase lies upstream of MAP kinases and stimulates the enzymatic activity of MAP kinases upon wide variety of extra- and intracellular signals. As an essential component of MAP kinase signal transduction pathway, this kinase is involved in many cellular processes such as proliferation, differentiation, transcription regulation and development. [provided by RefSeq, Jul 2008]

Immunogen

A synthetic peptide of human MEK1

Gene ID

5604

Swiss Prot

Q02750

Synonyms

CFC3; MEK1; MKK1; MAPKK1; PRKMK1

Reactivity

Human

Application

WB, IHC-P, ICC/IF, FC, IP

Calculated MW

43 kDa

Observed MW

43 kDa

Host species

Rabbit

Clonality

Monoclonal

Clonality No.

DGR11737


Isotype

IgG


Purity

Affinity Purification

Conjugation	Un-conjugated
Concentration	1 mg/mL
Formulation	PBS Only
Storage Stability	Store at -20°C. Recommended to aliquot into single-use vials. Supplied in 1X PBS (pH 7.4). BSA and Azide Free. Stable for 12 months from date of receipt.



Western blot analysis of extracts from Jurkat cells using [db15920](#) at 1:1000.



Immunofluorescence analysis of HeLa cells labelling MEK1 with [db15920](#).

The cells were fixed with 4% PFA (10min, RT) followed by treatment with 0.1% Triton X-100 (10min, RT), and blocked in 1% BSA/10% normal goat serum/0.3M glycine in 0.1% PBS-Tween 20 for 1h. The cells were then incubate with [db15920](#) (1:50) at room temprature for 1h, followed by a further incubation at room temperature for 45min with Goat Anti Rabbit IgG (H+L)-AF488 [db10005](#), shown in green). Nuclear DNA was labeled in blue with DAPI.

Control: Secondary antibody only.