
HDAC1 (3F5) Mouse mAb (PBS Only)

db6538-PBS **Package：可询价**


Product Name : HDAC1 (3F5) Mouse mAb (PBS Only)
Cat.No.: db6538-PBS
Synonyms : HDAC1; RPD3L1; Histone deacetylase 1; HD1
Application : IHC-P
Reactivity : Human, Rat, Mouse
Host species : Mouse

Background	Responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events. Histone deacetylases act via the formation of large multiprotein complexes. Deacetylates SP proteins, SP1 and SP3, and regulates their function. Component of the BRG1-RB1-HDAC1 complex, which negatively regulates the CREST-mediated transcription in resting neurons. Upon calcium stimulation, HDAC1 is released from the complex and CREBBP is recruited, which facilitates transcriptional activation. Deacetylates TSHZ3 and regulates its transcriptional repressor activity. Deacetylates 'Lys-310' in RELA and thereby inhibits the transcriptional activity of NF-kappa-B. Deacetylates NR1D2 and abrogates the effect of KAT5-mediated relieving of NR1D2 transcription repression activity. Component of a RCOR/GFI/KDM1A/HDAC complex that suppresses, via histone deacetylase (HDAC) recruitment, a number of genes implicated in multilineage blood cell development. Involved in CIART-mediated transcriptional repression of the circadian transcriptional activator/ CLOCK-ARNTL/BMAL1 heterodimer. Required for the transcriptional repression of circadian target genes, such as PER1, mediated by the large PER complex or CRY1 through histone deacetylation.
Immunogen	Synthetic peptide conjugated to KLH
Gene ID	3065
Swiss Prot	Q13547
Synonyms	HDAC1; RPD3L1; Histone deacetylase 1; HD1
Reactivity	Human, Rat, Mouse
Application	IHC-P
Calculated MW	62 kDa
Host species	Mouse

Clonality	Monoclonal
Clonality No.	3F5-2E8-1F7
Isotype	IgG1
Purity	Affinity Purification
Conjugation	Un-conjugated
Concentration	1 mg/mL
Formulation	PBS Only
Storage Stability	Store at -20°C. Recommended to aliquot into single-use vials. Supplied in 1X PBS (pH 7.4). BSA and Azide Free. Stable for 12 months from date of receipt.



Immunohistochemistry analysis of paraffin-embedded Human Colon Carcinoma Tissue using HDAC1 (3F5) antibody.High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.



Immunohistochemical analysis of paraffin-embedded Human tonsils using HDAC1 (3F5) antibody.High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.