

Cytochrome C (7C10) Mouse mAb (PBS Only)**db6489-PBS****Package :** 可询价**Product Name :** Cytochrome C (7C10) Mouse mAb (PBS Only)**Cat.No.:** db6489-PBS**Synonyms :** CYCS; CYC; Cytochrome c**Application :** WB, ICC/IF, IHC-P**Reactivity :** Human, Mouse, Rat, Chicken**Host species :** Mouse**Background**

Electron carrier protein. The oxidized form of the cytochrome c heme group can accept an electron from the heme group of the cytochrome c1 subunit of cytochrome reductase. Cytochrome c then transfers this electron to the cytochrome oxidase complex, the final protein carrier in the mitochondrial electron-transport chain. Plays a role in apoptosis. Suppression of the anti-apoptotic members or activation of the pro-apoptotic members of the Bcl-2 family leads to altered mitochondrial membrane permeability resulting in release of cytochrome c into the cytosol. Binding of cytochrome c to Apaf-1 triggers the activation of caspase-9, which then accelerates apoptosis by activating other caspases.

Immunogen

Recombinant Protein of CYCS

Gene ID

54205

Swiss Prot

P99999

Synonyms

CYCS; CYC; Cytochrome c

Reactivity

Human, Mouse, Rat, Chicken

Application

WB, ICC/IF, IHC-P

Calculated MW

12 kDa

Observed MW

12 kDa

Host species

Mouse

Clonality

Monoclonal

Clonality No.

7C10-3H3-9A3

Isotype

IgG1


Purity

Affinity Purification


Conjugation

Un-conjugated


Concentration	1 mg/mL
Formulation	PBS Only
Storage Stability	Store at -20°C. Recommended to aliquot into single-use vials. Supplied in 1X PBS (pH 7.4). BSA and Azide Free. Stable for 12 months from date of receipt.




Western blot analysis of Cytochrome c in T47D, 3T3, COS7, C6 and Hela lysates using Cytochrome c antibody.



Western blot analysis of Cytochrome C (7C10) in Hela, T47D, mouse brain, K562, rat brain lysates using Cytochrome C (7C10) antibody.



Immunocytochemistry analysis of Cytochrome C (7C1) (green) in Hela using Cytochrome C (7C1) antibody, and DAPI (blue)



Immunohistochemistry analysis of paraffin-embedded human tonsil tissue using Cytochrome C (7C10) antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.