

PYK2 (4B4) Mouse mAb (PBS Only)

db6398-PBS

Package : 可询价

Product Name : PYK2 (4B4) Mouse mAb (PBS Only)**Cat.No.:** db6398-PBS**Synonyms** : PKB; PTK; CAKB; FAK2; FRNK; PTK2B**Application** : WB, IHC-P**Reactivity** : Human**Host species** : Mouse**Background**

PYK2/ PTK2B protein tyrosine kinase 2 beta, also known as PTK2B, PKB, PTK, CAKB, FAK2, FRNK. Entrez Protein NP_004094. It is a cytoplasmic protein tyrosine kinase which is involved in calcium-induced regulation of ion channels and activation of the map kinase signaling pathway. The encoded protein may represent an important signaling intermediate between neuropeptide-activated receptors or neurotransmitters that increase calcium flux and the downstream signals that regulate neuronal activity. The encoded protein undergoes rapid tyrosine phosphorylation and activation in response to increases in the intracellular calcium concentration, nicotinic acetylcholine receptor activation, membrane depolarization, or protein kinase C activation. This protein has been shown to bind CRK-associated substrate, nephrocystin, GTPase regulator associated with FAK, and the SH2 domain of GRB2. The encoded protein is a member of the FAK subfamily of protein tyrosine kinases but lacks significant sequence similarity to kinases from other subfamilies. Four transcript variants encoding two different isoforms have been found for this gene.

Immunogen

Purified recombinant fragment of PYK2 (aa815-997) expressed in E. Coli

Gene ID

2185

Swiss Prot

Q14289

Synonyms

PKB; PTK; CAKB; FAK2; FRNK; PTK2B

Reactivity

Human

Application

WB, IHC-P

Calculated MW

116 kDa

Observed MW

116 kDa

Host species

Mouse


Clonality

Monoclonal


Clonality No.

4B4-D0-F3

Isotype	IgG2a
Purity	Affinity Purification
Conjugation	Un-conjugated
Concentration	1 mg/mL
Formulation	PBS Only
Storage Stability	Store at -20°C. Recommended to aliquot into single-use vials. Supplied in 1X PBS (pH 7.4). BSA and Azide Free. Stable for 12 months from date of receipt.



Western blot analysis of Pyk2 in A549, Ramos, Jurkat, THP-1 and Raji lysates using Pyk2 antibody.



Immunohistochemistry analysis of paraffin-embedded Human Tonsil tissues using PYK2 antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.