

Cyclin D2 (6E11) Mouse mAb (PBS Only)

db6075-PBS

Package : 可询价

Product Name : Cyclin D2 (6E11) Mouse mAb (PBS Only)**Cat.No.:** db6075-PBS**Synonyms** : KIAK0002**Application** : WB**Reactivity** : Human**Host species** : Mouse**Background**

The protein encoded by this gene belongs to the highly conserved cyclin family, whose members are characterized by a dramatic periodicity in protein abundance through the cell cycle. Cyclins function as regulators of CDK kinases. Different cyclins exhibit distinct expression and degradation patterns which contribute to the temporal coordination of each mitotic event. This cyclin forms a complex with CDK4 or CDK6 and functions as a regulatory subunit of the complex, whose activity is required for cell cycle G1/S transition. This protein has been shown to interact with and be involved in the phosphorylation of tumor suppressor protein Rb. Knockout studies of the homologous gene in mouse suggest the essential roles of this gene in ovarian granulosa and germ cell proliferation. High level expression of this gene was observed in ovarian and Ticular tumors. Mutations in this gene are associated with megalencephaly-polymicrogyria-polydactyly-hydrocephalus syndrome 3 (MPPH3).

Immunogen

Purified recombinant human Cyclin D2 protein fragments expressed in E.coli

Gene ID

894

Swiss Prot

P30279

Synonyms

KIAK0002

Reactivity

Human

Application

WB

Calculated MW

33 kDa

Observed MW

38 kDa

Host species

Mouse

Clonality

Monoclonal


Clonality No.

6E11-G6-F5

Isotype

IgG2b

Purity	Affinity Purification
Conjugation	Un-conjugated
Concentration	1 mg/mL
Formulation	PBS Only
Storage Stability	Store at -20°C. Recommended to aliquot into single-use vials. Supplied in 1X PBS (pH 7.4). BSA and Azide Free. Stable for 12 months from date of receipt.



Western blot analysis of Cyclin D2 in CHO-K1 lysates and CHO-K1 transfected by Cyclin D2 fragment EGFP fusion protein lysates using Cyclin D2 antibody.