

**NF- $\kappa$ B p65 (3D2) Mouse mAb (PBS Only)****db6456-PBS****Package :** 可询价**Product Name :** NF- $\kappa$ B p65 (3D2) Mouse mAb (PBS Only)**Cat.No.:** db6456-PBS**Synonyms :** NFKB3; RELA; TF65; Transcription factor p65; p65; NFkB**Application :** WB, IHC-Fr, IHC-P, ICC/IF, IP**Reactivity :** Human, Mouse, Rat**Host species :** Mouse**Background**

NF- $\kappa$ B is a pleiotropic transcription factor present in almost all cell types and is the endpoint of a series of signal transduction events that are initiated by a vast array of stimuli related to many biological processes such as inflammation, immunity, differentiation, cell growth, tumorigenesis and apoptosis. NF- $\kappa$ B is a homo- or heterodimeric complex formed by the Rel-like domain-containing proteins RELA/p65, RELB, NFKB1/p105, NFKB1/p50, REL and NFKB2/p52 and the heterodimeric p65-p50 complex appears to be most abundant one. The dimers bind at  $\kappa$ -B sites in the DNA of their target genes and the individual dimers have distinct preferences for different  $\kappa$ -B sites that they can bind with distinguishable affinity and specificity. Different dimer combinations act as transcriptional activators or repressors, respectively. NF- $\kappa$ B is controlled by various mechanisms of post-translational modification and subcellular compartmentalization as well as by interactions with other cofactors or corepressors. NF- $\kappa$ B complexes are held in the cytoplasm in an inactive state complexed with members of the NF- $\kappa$ B inhibitor ( $I$ - $\kappa$ B) family. In a conventional activation pathway,  $I$ - $\kappa$ B is phosphorylated by  $I$ - $\kappa$ B kinases (IKKs) in response to different activators, subsequently degraded thus liberating the active NF- $\kappa$ B complex which translocates to the nucleus. NF- $\kappa$ B heterodimeric p65-p50 and p65-c-Rel complexes are transcriptional activators. The NF- $\kappa$ B p65-p65 complex appears to be involved in invasin-mediated activation of IL-8 expression. The inhibitory effect of  $I$ - $\kappa$ B upon NF- $\kappa$ B in the cytoplasm is exerted primarily through the interaction with p65. p65 shows a weak DNA-binding site which could contribute directly to DNA binding in the NF- $\kappa$ B complex. Associates with chromatin at the NF- $\kappa$ B promoter region via association with DDX1. Essential for cytokine gene expression in T-cells (PubMed/15790681).

**Immunogen**

Recombinant Protein of Transcription factor p65

**Gene ID**

5970

**Swiss Prot**


Q04206

**Synonyms**


NFKB3; RELA; TF65; Transcription factor p65; p65; NFkB

---


<b>Reactivity</b>	Human, Mouse, Rat
<b>Application</b>	WB, IHC-Fr, IHC-P, ICC/IF, IP
<b>Calculated MW</b>	60 kDa
<b>Observed MW</b>	65 kDa
<b>Host species</b>	Mouse
<b>Clonality</b>	Monoclonal
<b>Clonality No.</b>	3D2-4E9-7A8
<b>Isotype</b>	IgG1
<b>Purity</b>	Affinity Purification
<b>Conjugation</b>	Un-conjugated
<b>Concentration</b>	1 mg/mL
<b>Formulation</b>	PBS Only
<b>Storage Stability</b>	Store at -20°C. Recommended to aliquot into single-use vials. Supplied in 1X PBS (pH 7.4). BSA and Azide Free. Stable for 12 months from date of receipt.




Western blot analysis of NF-KB p65 in T47D, 3T3, COS7, C6 and HeLa lysates using NF-KB p65 antibody.




Immunofluorescence analysis of NF-KB p65 (3D2) in HeLa using NF-KB p65 (3D2) antibody, and DAPI (blue).



Immunoprecipitation analysis of NF-KB p65 (3D2) in HeLa lysates using NF-KB p65 (3D2) antibody.



Immunohistochemistry analysis of paraffin-embedded rat Brain Tissue using NF-KB p65 antibody. High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.



Western blot analysis of NF-KB p65 (3D2) in COS7, C6, T47D lysates using NF-KB p65 (3D2) antibody.