
NF-kB p65 (3D2) Mouse mAb (PBS Only)

db6456-PBS **Package：可询价**


Product Name : NF-kB p65 (3D2) Mouse mAb (PBS Only)
Cat.No.: db6456-PBS
Synonyms : NFKB3; RELA; TF65; Transcription factor p65; p65; NFkB
Application : WB, IHC-Fr, IHC-P, ICC/IF, IP
Reactivity : Human, Mouse, Rat
Host species : Mouse

Background	NF-kappa-B is a pleiotropic transcription factor present in almost all cell types and is the endpoint of a series of signal transduction events that are initiated by a vast array of stimuli related to many biological processes such as inflammation, immunity, differentiation, cell growth, tumorigenesis and apoptosis. NF-kappa-B is a homo- or heterodimeric complex formed by the Rel-like domain-containing proteins RELA/p65, RELB, NFKB1/p105, NFKB1/p50, REL and NFKB2/p52 and the heterodimeric p65-p50 complex appears to be most abundant one. The dimers bind at kappa-B sites in the DNA of their target genes and the individual dimers have distinct preferences for different kappa-B sites that they can bind with distinguishable affinity and specificity. Different dimer combinations act as transcriptional activators or repressors, respectively. NF-kappa-B is controlled by various mechanisms of post-translational modification and subcellular compartmentalization as well as by interactions with other cofactors or corepressors. NF-kappa-B complexes are held in the cytoplasm in an inactive state complexed with members of the NF-kappa-B inhibitor (I-kappa-B) family. In a conventional activation pathway, I-kappa-B is phosphorylated by I-kappa-B kinases (IKKs) in response to different activators, subsequently degraded thus liberating the active NF-kappa-B complex which translocates to the nucleus. NF-kappa-B heterodimeric p65-p50 and p65-c-Rel complexes are transcriptional activators. The NF-kappa-B p65-p65 complex appears to be involved in invasin-mediated activation of IL-8 expression. The inhibitory effect of I-kappa-B upon NF-kappa-B the cytoplasm is exerted primarily through the interaction with p65. p65 shows a weak DNA-binding site which could contribute directly to DNA binding in the NF-kappa-B complex. Associates with chromatin at the NF-kappa-B promoter region via association with DDX1. Essential for cytokine gene expression in T-cells (PubMed/15790681).
Immunogen	Recombinant Protein of Transcription factor p65
Gene ID	5970
Swiss Prot	Q04206
Synonyms	NFKB3; RELA; TF65; Transcription factor p65; p65; NFkB


Reactivity	Human, Mouse, Rat
Application	WB, IHC-Fr, IHC-P, ICC/IF, IP
Calculated MW	60 kDa
Observed MW	65 kDa
Host species	Mouse
Clonality	Monoclonal
Clonality No.	3D2-4E9-7A8
Isotype	IgG1
Purity	Affinity Purification
Conjugation	Un-conjugated
Concentration	1 mg/mL
Formulation	PBS Only
Storage Stability	Store at -20°C. Recommended to aliquot into single-use vials. Supplied in 1X PBS (pH 7.4). BSA and Azide Free. Stable for 12 months from date of receipt.




Western blot analysis of NF-KB p65 in T47D, 3T3, COS7, C6 and Hela lysates using NF-KB p65 antibody.




Immunofluorescence analysis of NF-KB p65 (3D2) in Hela using NF-KB p65 (3D2) antibody,and DAPI (blue).



Immunoprecipitation analysis of NF-KB p65 (3D2) in Hela lysates using NFκB p65 (3D2) antibody.



Immunohistochemistry analysis of paraffin-embedded rat Brain Tissue using NF-KB p65 antibody.High-pressure and temperature Sodium Citrate pH 6.0 was used for antigen retrieval.



Western blot analysis of NF-KB p65 (3D2) in COS7, C6, T47D lysates using NF-KB p65 (3D2) antibody.